

PCT

WORLD INTELLECTUAL PROPERTY
ORGANIZATION

INTERNATIONAL APPLICATION PUBLISHED UNDER 1

WO 9605188A1

(51) International Patent Classification: **C07D 281/00, A13J04, A61K 31/55**

(11) International Publication Number: **WO 9605188**

(43) International Publication Date: **22 February 1996 (22.02.96)**

(21) International Application Number: **PCT/GB95/01884**

(22) International Filing Date: **9 August 1995 (09.08.95)**

(30) Priority Date: **08/288,577** **10 August 1994 (10.08.94)** **US**

(71) Applicant (for all designated States except US): **THE WELL-COME FOUNDATION LIMITED (GB)**; **Unicorn House, 160 Euston Road, London NW1 2BP (GB).**

(72) Inventor and Inventor/Applicant (for US only): **BREADDY, Lawrence, Edward (US/US); 5924 Wintergreen Drive, Raleigh, NC 27609 (US).**


(74) Agent: **McLEOD HAMMER, Carltons; The Wellcome Foundation Limited, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).**

Published
With international search report.

(54) Title: **HYPOLIPIDEMIC 1,4-BENZOTHAZEPINE-1,1-DIOXIDES**

(57) Abstract

The invention is concerned with novel hypolipidemic compounds of formula (I), with processes and novel intermediates for their preparation, pharmaceutical compositions containing them and with their use in medicine, particularly in the prophylaxis and treatment of hyperlipidemic conditions, such as atherosclerosis.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Morocco
AU	Australia	GE	Georgia	MT	Malta
BA	Bosnia	GR	Greece	NL	Netherlands
BB	Barbados	HN	Honduras	NO	Norway
BE	Belgium	HU	Hungary	NZ	New Zealand
BF	Burkina Faso	IE	Ireland	PL	Poland
BG	Bulgaria	IT	Italy	PT	Portugal
BR	Brazil	JP	Japan	RU	Russia
BS	Bahamas	KE	Kenya	SD	Sudan
CA	Canada	KG	Kyrgyzstan	TD	Chad
CC	Cocos (Keeling) Islands	KZ	Kazakhstan	TG	Togo
CF	Congo	LA	Laos	TT	Trinidad and Tobago
CG	Congo (Brazzaville)	LV	Latvia	UA	Ukraine
CH	Switzerland	MC	Monaco	US	United States of America
CI	Cote d'Ivoire	MD	Moldova	UZ	Uzbekistan
CM	Cameroun	MG	Madagascar	VN	Viet Nam
CN	China	MN	Mongolia		
CO	Colombia				
CZ	Czech Republic				
DE	Germany				
DK	Denmark				
DM	Dominica				
DO	Dominican Republic				
EE	Estonia				
EG	Egypt				
ES	Spain				
FI	Finland				
FR	France				
GA	Gabon				

HYPOLIPIDEMIC 1,4-BENZOTHAZEPINE-1,1-DIOXIDES

1


The present invention is concerned with new hypolipidemic compounds, with processes and novel intermediates for their preparation, with pharmaceutical compositions containing them and with their use in medicine, particularly in the prophylaxis and treatment of hyperlipidemic conditions, such as atherosclerosis.

Hyperlipidemic conditions are often associated with elevated plasma concentrations of low density lipoprotein (LDL) cholesterol and very low density lipoprotein (VLDL) cholesterol. Such concentrations can be reduced by decreasing the absorption of bile acids from the intestine. One method by which this may be achieved is to inhibit the bile acid uptake system in the terminal ileum. Such inhibition stimulates the conversion of cholesterol to bile acid by the liver and the resulting increase in demand for cholesterol produces a corresponding increase in the rate of clearance of LDL and VLDL cholesterol from the blood plasma or serum.

There has now been identified a novel class of heterocyclic compounds which reduce the plasma or serum concentrations of LDL and VLDL cholesterol and in consequence are particularly useful as hypolipidemic agents. By decreasing the concentrations of cholesterol and cholesterol ester in the plasma, the compounds of the present invention retard the build-up of atherosclerotic lesions and reduce the incidence of coronary heart disease-related events. The latter are defined as cardiac events associated with increased concentrations of cholesterol and cholesterol ester in the plasma or serum.

For the purposes of this specification, a hyperlipidemic condition is defined as any condition wherein the total cholesterol concentration (LDL + VLDL) in the plasma or serum is greater than 240mg/dL (6.21mmol/L) (J. Amer. Med. Assn. 256, 20, 2849-2858 (1986)).

International Patent Application No. WO 93/16055 describes compounds of formula (O)



2

wherein

l is an integer of from 0 to 4;

m is an integer of from 0 to 5;

n is an integer of from 0 to 2;


R and R' are atoms or groups independently selected from halogen, nitro, phenylalkoxy, C₁₋₄ alkoxy, C₁₋₆ alkyl, and -O(CH₂)_pSO₂R'' wherein p is an integer of from 1 to 4 and R'' is hydrogen or C₁₋₆ alkyl, wherein said phenylalkoxy, alkoxy and alkyl groups are optionally substituted by one or more halogen atoms;

R^A is a C₁₋₆ straight, that is, unbranched, alkyl group, and


R^B is a C₂₋₆ straight, that is, unbranched, alkyl group, and salts, solvates and physiologically functional derivatives thereof, as being useful as hypolipidemic agents.

We have now discovered a group of compounds which have greater hypolipidemic activity in vivo than those specifically disclosed in International Patent Application No. 93/16055.

Accordingly, the present invention provides compounds of the formula (I):



wherein R¹ is a straight chained C₁₋₆ alkyl group, R² is a straight chained C₁₋₆ alkyl group, R³ is hydrogen or a group OR¹¹ in which R¹¹ is hydrogen, optionally substituted C₁₋₆ alkyl or a C₁₋₆ alkylcarbonyl group, R⁴ is pyridyl or optionally substituted phenyl, R⁵, R⁶, R⁷ and R⁸ are the same or different and each is selected from hydrogen, halogen, cyano, R¹⁵-acylide, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, P(O)(OR¹⁵)₂, OCO¹⁵, OCF₃, OCN, SCN, NHCN, CH₃OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO¹⁵, (CH₂)_pNR¹²R¹³, CO¹⁵R¹⁵, NHCO¹⁵, NHSO¹⁵, R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pSO₂R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein p is an integer from 1-4, n is an integer from 0-3 and R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from hydrogen and optionally substituted C₁₋₆ alkyl; or R⁶ and R⁷ are linked to form a group



wherein R¹² and R¹³ are as hereinbefore defined and m is 1 or 2, and R⁹ and R¹⁰ are the same or different and each is hydrogen or C₁₋₆ alkyl; and salts, solvates or a physiologically functional derivatives thereof, with the proviso that when R³ is hydrogen either R⁷ is not hydrogen or at least two of R⁵, R⁶, R⁷ and R⁸ are not hydrogen; and salts, solvates, and physiologically functional derivatives thereof.

When R⁴ is a substituted phenyl group, there may be one to five, preferably one or two substituents which are the same or different and are each selected from halogen, hydroxy, nitro, phenyl-C₁₋₆ alkoxy, C₁₋₆ alkoxy, optionally substituted C₁₋₆ alkyl, S(O)_nR¹⁵, CO₂R¹⁵, O(CH₂CH₂O)_nR¹⁵, O(CH₂)_pSO₃R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein R¹² to R¹⁵, n and p are as hereinbefore defined.

According to a further aspect, the invention provides compounds of formula (I) wherein:

R¹ and R² are straight chained C₁₋₆ alkyl;

R³ is hydrogen or hydroxy;

R⁴ is unsubstituted phenyl;

R⁵ is hydrogen;

R⁹ and R¹⁰ are both hydrogen; and either

R⁷ is selected from halogen, hydroxy, C₁₋₆ alkoxy, optionally substituted C₁₋₆ alkyl, -S(O)_nR¹⁵, -OC(O)R¹⁵, and -CH₂OR¹⁵ wherein R¹⁵ is hydrogen or C₁₋₆ alkyl; and





R⁶ and R⁸ are independently selected from hydrogen and those groups listed in the definition of R⁷; or

R⁸ is hydrogen and R⁶ and R⁷ are linked to form a group -O-(CH₂)_m-O- wherein m is 1 or 2;

and salts, solvates, and physiologically functional derivatives thereof

Of the compounds of formula (I), those in which R⁸ is hydrogen and R⁶ and R⁷ are both C₁₋₆ alkoxy, more particularly both methoxy, are preferred.

Preferred embodiments of the compounds of formula (I) include compounds of the formula (II), (III), (IV) or (IVa)



wherein R¹ to R¹⁰ are as hereinbefore defined and R^{7a} is selected from halogen, cyano, R¹⁵-acetylide, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, P(O)(OR¹⁵)₂, OCOR¹⁵, OCF₃, OCN, SCN, HNCN, CH₂OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHSO₂R¹⁵, OCH₂OR¹⁵, OCH=CH-R¹⁵, O(CH₂CH₂O)_pR¹⁵, O(CH₂)_pSO₃R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein n, p and R¹² to R¹⁵ are as hereinbefore defined; with the proviso that in compounds of formula (III) at least two of R⁵ to R⁸ are not hydrogen; and salts solvates and physiologically functional derivatives thereof.

When one or more of R³ to R⁸ or R¹¹ to R¹⁴ is a substituted C₁₋₆ alkyl group, or comprises a C₁₋₆ alkyl group the substituents may be the same or different and each is selected from hydroxy, halogen, C₁₋₆ alkoxy, COR¹⁶, nitrile, CO₂R¹⁶,

SO_3R^{16} , $NR^{17}R^{18}$, $N^+R^{17}R^{18}R^{19}$ wherein R^{16} to R^{19} are the same or different and each is selected from hydrogen or C_{1-6} alkyl.

Suitably R^1 is methyl, ethyl or n-propyl and preferably R^1 is ethyl. Suitably R^2 is methyl, ethyl, n-propyl, n-butyl or n-pentyl. Preferably R^2 is n-butyl.

Preferably R^5 is hydrogen.

Suitably R^7 and R^{7a} are selected from OR^{15} , SO_2R^{15} , $OCOR^{15}$, OCE_3 , OCN , SCN , CHO , OCH_2OR^{15} , $OCH=CHR^{15}$, $O(CH_2CH_2O)_pR^{15}$, $O(CH_2)_pSO_3R^{15}$, $O(CH_2)_pNR^{12}R^{13}$ and $O(CH_2)_pN^+R^{12}R^{13}R^{14}$ wherein p is an integer from 1-4, n is an integer from 0-3 and R^{12} , R^{13} , R^{14} and R^{15} are independently selected from hydrogen and optionally substituted C_{1-6} alkyl. Preferably R^7 and R^{7a} are OR^{15} .

Suitably R^9 and R^{10} are hydrogen, methyl or ethyl. Preferably R^9 and R^{10} are both hydrogen.

Suitably R^4 is pyridyl or phenyl optionally substituted, preferably at the 4- and/or 3-position by halogen, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, hydroxy, carboxy or $O(CH_2)_3SO_3H$. Preferably R^4 is unsubstituted phenyl.

In the compounds of the formula (II) : suitably one or two, and preferably one, of R^5 , R^6 and R^8 is other than hydrogen and suitably each is selected from C_{1-4} alkyl, optionally substituted by fluoro, C_{1-4} alkoxy, halogen and hydroxy. Most suitably, each is selected from methyl, methoxy, hydroxy, trifluoromethyl and halo. Preferably, R^6 is methoxy or bromo and R^5 and R^8 are hydrogen. Suitably, R^{7a} is C_{1-4} alkyl optionally substituted by fluoro, C_{1-4} alkoxy, halogen or hydroxy. Most suitably, R^{7a} is methoxy, hydroxy or trifluoromethyl and preferably R^{7a} is methoxy.

In the compounds of the formula (III) : suitably at least one and preferably two of R^5 to R^8 are hydrogen. Preferably at least one of R^6 and R^7 is not hydrogen. When R^5 to R^8 are other than hydrogen then they are suitably C_{1-4} alkyl optionally substituted by fluoro, C_{1-4} alkoxy, halogen or hydroxy, most suitably methyl, methoxy, hydroxy, trifluoromethyl or chloro and preferably methoxy.

In the compounds of the formula (IV) : suitably two, three or four of R^5 to R^8 are hydrogen, the others being C_{1-4} alkyl optionally substituted by fluoro, C_{1-4} alkoxy,

halogen or hydroxy and most suitably methyl, methoxy, hydroxy, trifluoromethyl or chloro and preferably methoxy.

In the compounds of formula (IVa) : suitably at least one and preferably two of R^5 to R^8 are hydrogen. Preferably at least one of R^6 and R^7 is not hydrogen. When R^5 to R^8 are other than hydrogen then they are suitably C_{1-4} alkyl optionally substituted by fluoro, C_{1-4} alkoxy, halogen or hydroxy, most suitably methyl, methoxy, hydroxy, trifluoromethyl or chloro and preferably methoxy. Most preferably, R^1 is n-butyl, R^2 is ethyl, R^3 , R^5 , R^8 , R^9 and R^{10} are hydrogen, R^4 is pyridyl or optionally substituted phenyl and R^6 and R^7 are methoxy.

Preferred compounds of formula (I) are selected from the group consisting of:

(3R,5R)-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide;

(3R,5R)-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-4-ol,1,1-dioxide;

(+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide;

(+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-4-ol,1,1-dioxide;

(3R,5R)-7-Bromo-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(3R,5R)-7-Bromo-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-4-ol,1,1-dioxide;

(3R,5R)-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-7,8-diol,1,1-dioxide;

(3R,5R)-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-ol,1,1-dioxide;

- (3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7-methoxy-5-phenyl-1,4-benzothiazepine-8-ol 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-ol 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-4,8-diol;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide;
- (+)-Trans-2-((3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepin-7-yl)methoxy) ethanol S,S-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-hydroxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-thiol 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-sulfonic acid 1,1-dioxide;
- (7R,9R)-7-Butyl-7-ethyl-6,7,8,9-tetrahydro-9-phenyl-1,3-dioxolane(4,5-H)(1,4)-benzothiazepine 5,5-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8,9-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (3R,5R)-3-Butyl-3-ethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepine-4-ol 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-methanol S,S-dioxide;

- (3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-7-nitro-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-7-(methoxymethyl)-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-7,8-diyl diacetate 1,1-dioxide;
- (8R,10R)-8-Butyl-8-ethyl-2,3,7,8,9,10-hexahydro-10-1,4-dioxono(2,3-H)(1,4)-benzothiazepine 6,6-dioxide;
- (3R,5R)-3-butyl-7,8-diethoxy-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-butyl-8-ethoxy-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-1,1-dioxide;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-isopropoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide hydrochloride;
- (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-carbaldehyde 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- 3,3-Diethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-8-methoxy-1,4-benzothiazepine 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (RS)-3,3-Diethyl-2,3,4,5-tetrahydro-4-hydroxy-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-butyl-8-ethoxy-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-4-ol 1,1-dioxide;

- (+)-Trans-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-8-isopropoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;
- (+)-Trans-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-7,8,9-trimethoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;
- (3R,5R)-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-4,7,8-triol 1,1-dioxide;
- (+)-Trans-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-4,7,8-trimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-benzyl-3-ethyl-5-phenyl-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepin-4-yl acetate S,S-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-7-methoxy-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;
- 3,3-Dibenzyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;
- (+)-Trans-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl hydrogen sulphate;
- (+)-Trans-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl dihydrogen phosphate;
- 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl hydrogen sulphate;
- (+)-Trans-3-benzyl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl aspartate; and

3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl aspartate.

Pharmaceutically acceptable salts are particularly suitable for medical applications because of their greater aqueous solubility relative to the parent, i.e. basic, compounds. Such salts must clearly have a pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the present invention include those derived from inorganic acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric, sulphuric and sulphuric acids, and organic acids, such as acetic, benzenesulphonic, benzoic, citric, ethanesulphonic, fumaric, gluconic, glycolic, isothionic, lactic, lactobionic, malic, malic, methanesulphonic, succinic, p-toluenesulphonic, tartaric and trifluoroacetic acids. The chloride salt is particularly preferred for medical purposes. Suitable pharmaceutically acceptable base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, and alkaline earth salts, such as magnesium and calcium salts.

Salts having a non-pharmaceutically acceptable anion are within the scope of the invention as useful intermediates for the preparation or purification of pharmaceutically acceptable salts and/or for use in non-therapeutic, for example, *in vitro*, applications.

The term "physiologically functional derivative" as used herein refers to any physiologically acceptable derivative of a compound of the present invention, for example, an ester, which upon administration to a mammal, such as a human, is capable of providing (directly or indirectly) such a compound or an active metabolite thereof.

A further aspect of the present invention is products of the compounds of the invention. Such products can be metabolised *in vivo* to give a compound according to the invention. These products may or may not be active in their own right.

The compounds of the present invention can also exist in different polymorphic forms, for example, amorphous and crystalline polymorphic forms. All polymorphic forms of the compounds of the present invention are within the scope of the invention and are a further aspect thereof.

The term "alkyl" as used herein refers, unless otherwise stated, to a monovalent straight or branched chain radical. Likewise, the term "alkoxy" refers to a monovalent straight or branched chain radical attached to the parent molecular moiety through an oxygen atom. The term "phenylalkoxy" refers to a monovalent phenyl group attached to a divalent C₁₋₆

alkylene group which is itself attached to the parent molecular moiety through an oxygen atom.

The compounds of formula (I) exist in forms wherein the carbon centres $-C(R^1)(R^2)-$ and $-CHR^4-$ is/are chiral. The present invention includes within its scope each possible optical isomer substantially free, i.e. as associated with less than 5%, of any other optical isomer(s), and mixtures of one or more optical isomers in any proportions, including racemic mixtures.

For the purposes of this specification, the absolute chiralities of the aforementioned carbon centres are given in the order $-C(R^1)(R^2)-$, then $-CHR^4-$.

In those cases where the absolute stereochemistry at $-C(R^1)(R^2)-$ and $-CHR^4-$ has not been determined, the compounds of the invention are defined in terms of the relative positions of the R^1/R^2 and H/R^4 substituents. Thus those compounds wherein the bulkier of the R^1 and R^2 substituents, i.e. the substituent of higher mass, and the R^4 substituent are both located on the same side of the thiapine ring are referred to herein as "cis", and those compounds in which the bulkier of the R^1 and R^2 substituents are located on opposite sides of the ring are referred to as "trans" and are preferred. It will be evident to a skilled person that both "cis" and "trans" compounds of the invention can each exist in two enantiomeric forms which are individually designated "(+)" or "(-)" according to the direction of rotation of a plane of polarised light when passed through a sample of the compound. Cis or trans compounds of the invention in which the individual enantiomers have not been resolved are referred to herein using the prefix "(+/-)".

According to further aspects of the invention, there are also provided:

- (a) compounds of formula (I) and pharmaceutically acceptable salts, solvates and physiologically functional derivatives thereof for use as therapeutic agents, particularly in the prophylaxis and treatment of clinical conditions for which a bile acid uptake inhibitor is indicated, for example, a hyperlipidemic condition, such as atherosclerosis;
- (b) pharmaceutical compositions comprising a compound of formula (I) or one of its pharmaceutically acceptable salts, solvates, or physiologically functional derivatives, at least one pharmaceutically acceptable carrier and, optionally, one or more other physiologically active agents;

- (c) the use of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof in the manufacture of a medicament for the prophylaxis or treatment of a clinical condition for which a bile acid uptake inhibitor is indicated, for example, a hyperlipidemic condition, such as atherosclerosis;
- (d) a method of inhibiting the absorption of bile acids from the intestine of a mammal, such as a human, which comprises administering an effective bile acid absorption inhibiting amount of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof to the mammal;
- (e) a method of reducing the blood plasma or serum concentrations of LDL and VLDL cholesterol in a mammal, such as a human, which comprises administering an effective cholesterol reducing amount of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof to the mammal;
- (f) a method of reducing the concentrations of cholesterol and cholesterol ester in the blood plasma or serum of a mammal, such as a human, which comprises administering an effective cholesterol and cholesterol ester reducing amount of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof to the mammal;
- (g) a method of increasing the fecal excretion of bile acids in a mammal, such as a human, which comprises administering an effective bile acid fecal excretion increasing amount of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof to the mammal;
- (h) a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a bile acid uptake inhibitor is indicated, for example, a hyperlipidemic condition, such as atherosclerosis, which comprises administering a therapeutically effective amount of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof to the mammal;
- (i) a method of reducing the incidence of coronary heart disease-related events in a mammal, such as a human, which comprises administering an effective coronary heart disease-related events reducing amount of a compound of formula (I) or of a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof;

13

- (j) a method of reducing the concentration of cholesterol in the blood plasma or serum of a mammal, such as a human, which comprises administering an effective cholesterol reducing amount of a compound of formula (I);
- (k) processes for the preparation of compounds of formula (I) (including salts, solvates and physiologically functional derivatives thereof as defined herein); and
- (l) novel chemical intermediates in the preparation of compounds of formula (I).
- (m) the compounds of Synthetic Examples 1-53 as hereinafter disclosed.

Hereinafter all references to "compound(s) of formula (I)" refer to compound(s) of formula (I) as described above together with their salts, solvates and physiologically functional derivatives as defined herein.

The amount of a compound of formula (I) which is required to achieve the desired biological effect will, of course, depend on a number of factors, for example, the specific compound chosen, the use for which it is intended, the mode of administration and the clinical condition of the recipient. In general, a daily dose is in the range of from 0.3mg to 100mg (typically from 3mg to 50mg) per day per kilogram bodyweight, for example, 3-10mg/kg/day. An intravenous dose can, for example, be in the range of from 0.3mg to 1.0mg/kg, which can conveniently be administered as an infusion of from 10mg to 100mg per kilogram per minute. Infusion fluids suitable for this purpose can contain, for example, from 0.1mg to 10mg, typically from 1mg to 10mg, per millilitre. Unit doses can contain, for example, from 1mg to 10g of the active compound. Thus ampoules for injection can contain, for example, from 1mg to 100mg and orally administrable unit dose formulations, such as tablets or capsules, may contain, for example, from 1.0 to 1000mg, typically from 10 to 600mg. In the case of pharmaceutically acceptable salts, the weights indicated above refer to the weight of the benzothiazepine ion derived from the salt.

For the prophylaxis or treatment of the conditions referred to above, the compounds of formula (I) can be used as the compound *per se*, but are preferably presented with an acceptable carrier in the form of a pharmaceutical composition. The carrier must, of course, be acceptable in the sense of being compatible with the other ingredients of the composition and must not be deleterious to the recipient. The carrier can be a solid or a liquid, or both, and is preferably formulated with the compound as a unit-dose composition, for example, a tablet, which can contain from 0.05% to 95% by weight of the active compound. Other

14

pharmacologically active substances can also be present including other compounds of formula (I). The pharmaceutical compositions of the invention can be prepared by any of the well known techniques of pharmacy consisting essentially of admixing the components.

Pharmaceutical compositions according to the present invention include those suitable for oral, rectal, topical, buccal (e.g. sub-lingual) and parenteral (e.g. subcutaneous, intramuscular, intradermal, or intravenous) administration, although the most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular compound of formula (I) which is being used. Enteric-coated and enteric-coated controlled release formulations are also within the scope of the invention. Suitable enteric coatings include cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate and anionic polymers of methacrylic acid and methacrylic acid methyl ester.

Pharmaceutical compositions suitable for oral administration can be presented in discrete units, such as capsules, cachets, lozenges, or tablets, each containing a predetermined amount of a compound of formula (I), as a powder or granules, as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil emulsion. As indicated, such compositions can be prepared by any suitable method of pharmacy which includes the step of bringing into association the active compound and the carrier (which can constitute one or more accessory ingredients). In general, the compositions are prepared by uniformly and intimately admixing the active compound with a liquid or finely divided solid carrier, or both, and then, if necessary, shaping the product. For example, a tablet can be prepared by compressing or moulding a powder or granules of the compound, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing, in a suitable machine, the compound in a free-flowing form, such as a powder or granules optionally mixed with a binder, lubricant, inert diluent and/or surface active/dispersing agent(s). Moulded tablets can be made by moulding, in a suitable machine, the powdered compound moistened with an inert liquid diluent.

Pharmaceutical compositions suitable for buccal (sub-lingual) administration include lozenges comprising a compound of formula (I) in a flavored base, usually sucrose and, acacia or tragacanth, and pastilles comprising the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

Pharmaceutical compositions suitable for parenteral administration conveniently comprise sterile aqueous preparations of a compound of formula (I), preferably isotonic with the

15

blood of the intended recipient. These preparations are preferably administered intravenously, although administration can also be effected by means of subcutaneous, intramuscular, or intradermal injection. Such preparations can conveniently be prepared by admixing the compound with water and rendering the resulting solution sterile and isotonic with the blood. Injectable compositions according to the invention will generally contain from 0.1 to 5% w/w of the active compound.

Pharmaceutical compositions suitable for rectal administration are preferably presented as unit-dose suppositories. These can be prepared by admixing a compound of formula (I) with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.


Pharmaceutical compositions suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, aerosol, or oil. Carriers which can be used include vaseline, lanoline, polyethylene glycols, alcohols, and combinations of two or more thereof. The active compound is generally present at a concentration of from 0.1 to 15% w/w of the composition, for example, from 0.5 to 2%.

Transdermal administration is also possible. Pharmaceutical compositions suitable for transdermal administration can be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Such patches suitably contain the active compound in an optionally buffered, aqueous solution, dissolved and/or dispersed in an adhesive, or dispersed in a polymer. A suitable concentration of the active compound is about 1% to 35%, preferably about 3% to 15%. As one particular possibility, the active compound can be delivered from the patch by electrotransport or iontophoresis, for example, as described in Pharmaceutical Research, 2(6), 318 (1986).

The compounds of the invention can be prepared by conventional methods known to a skilled person or in an analogous manner to processes described in the art.


For example, compounds of the formula (I) wherein R³ is hydrogen can be prepared by oxidation of the corresponding compound of the formula (V):

16



wherein R¹ to R¹⁰ are as hereinbefore defined and l is 0 or 1. This oxidation may suitably be carried out by reaction with a peroxide, for example hydrogen peroxide in the presence of trifluoroacetic acid at a non-extreme temperature, e.g. -20°C to 50°C and preferably at -10°C to 10°C. The compound of the formula (V) where l is 1 may be prepared from the corresponding compound where l is 0 by partial oxidation using a peroxide as described above.


Compounds of formula (V) can be prepared by reducing the imine bond of a compound of formula (VI)



wherein R¹ to R¹⁰ are as hereinbefore defined, using, for example, a boron compound, such as borane, in a suitable solvent, for example an ether such as THF, or catalytic hydrogenation using, for example, a palladium catalyst, such as 10% Pd/C at a non-extreme temperature, for example -20°C to 100°C and preferably -10°C to 50°C.


Compounds of formula (VI) as herein defined as well as each possible optical isomer substantially free, i.e., associated with less than 5% of any other optical isomer(s), and mixtures of one or more optical isomers in any proportions, including racemic mixtures are considered to be novel and constitute a further aspect of the present invention.

Compounds of formula (VI) can be prepared by cyclising compounds of formula (VII)




wherein R¹ to R⁸ are as hereinbefore defined, by, for example, azeotropic distillation or refluxing in the presence of a suitable drying agent, such as molecular sieves, in a suitable solvent, for example, 2,5-hexidine, in the presence of an acid, such as HCl.

Compounds of formula (VII) can be prepared by reacting a compound of formula (VIII)



wherein R⁴ to R⁸ are as hereinbefore defined, with the appropriately substituted aziridine, typically in a polar solvent, for example, methanol.

Compounds of formula (VII) can also be prepared by reacting a compound of formula (IX)




wherein R⁴ to R⁸ are as hereinbefore defined and L is a suitable leaving group, for example, halogen, with a compound of formula HSC(R⁹)(R¹⁰)C(R¹)(R²)NH₂, wherein R¹, R², R⁹ and R¹⁰ are as hereinbefore defined.

18

Compounds of formula (IX) can be prepared by reacting the corresponding acid with a compound of formula R⁴H wherein R⁴ is as hereinbefore defined, typically by a Friedel-Crafts reaction using, for example, aluminium chloride.

Compounds of formula (VIII) can be prepared by reacting a compound of formula (X)



wherein R⁵ to R⁸ are as hereinbefore defined with a non-nucleophilic base such as sodium hydride followed by treatment of the resulting salt with N,N-dimethylthiocarbonyl chloride, pyrolysis of the resulting O-aryldialkylthiocarbamate to the S-aryldialkylthiocarbamate (for example in a high boiling solvent such as tetradecane at a temperature of about 235°C.), and hydrolysis (for example with a strong base such as KOH).

Alternatively, compounds of formula (VIII) can be prepared by reacting compounds of formula (IX) with sodium hydrosulfide (NaSH).

The starting materials as hereinbefore defined can be obtained commercially or prepared by methods known to those skilled in the art or obtainable from the chemical literature, for example, the aziridines can be prepared from the corresponding 2-substituted 2-aminoethanols.

Compounds of the formula (V) wherein one or more of R⁵ to R⁸ is halo may be converted to compounds of the formula (V) wherein R⁵ to R⁸ is a different functional group by methods known to those skilled in the art or readily available from the literature.

Compounds of formula (I) wherein R³ = OH can be prepared from the corresponding compounds of formula (I) wherein R³ = H by oxidation with, for example, m-chloroperoxybenzoic acid.

The compounds of formula (I) substantially free, of other optical isomers can be obtained either by chiral synthesis, for example, by the use of the appropriate chiral starting

material(s), such as the aziridine, or by resolution of the products obtained from achiral syntheses, for example, by chiral hplc or by classical resolution with chiral acids.

Optional conversion of a compound of formula (I), or a compound of formula (I) comprising a basic substituent, to a corresponding acid addition salt may be effected by reaction with a solution of the appropriate acid, for example, one of those recited earlier. Optional conversion of a compound of formula (I) comprising an acidic substituent to a corresponding base salt may be effected by reaction with a solution of the appropriate base, for example, sodium hydroxide. Optional conversion to a physiologically functional derivative, such as an ester, can be carried out by methods known to those skilled in the art or obtainable from the chemical literature.

In addition, compounds of the formula (I) may be converted to different compounds of the formula (I) by standard methods known or available from the literature to those skilled in the art, for example by alkylation of a hydroxy group.

For a better understanding of the invention, the following Examples are given by way of illustration and are not to be construed in any way as limiting the scope of the invention.

Synthetic Example 1

Preparation of (R)-3R-3-butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide

(a) Ethyl 2-aminobutyrate hydrochloride

A slurry of 2-aminobutyric acid (100g, Aldrich) in absolute ethanol (300 ml) was stirred under nitrogen at 0°C and thionyl chloride (120.8g) was added dropwise. The reaction was stirred overnight at 0°C and then gradually warmed to room temperature. The resulting white slurry was heated under reflux for 3 hours, left to cool for 10 minutes, then poured into chilled diethyl ether (600ml), with hand stirring. The suspension was filtered and the solid product dried to give the desired product (150g) as a white solid. ¹H NMR consistent with proposed structure.

(b) Ethyl 2-benzylideneamino-2-ethylhexanoate

A solution of the product from step (a) (149.6g), magnesium sulfate (74.3g), and triethylamine (246 ml) in dichloromethane (1500 ml) was stirred at room temperature under

nitrogen and benzaldehyde (94.9g, Aldrich) was added dropwise. The mixture was stirred at room temperature for 3 hours then filtered. The filtrate was concentrated, triturated in diethyl ether, filtered and concentrated to yield the desired product as a yellow oil (174g). ¹H NMR consistent with the proposed structure.

(c) (±)-Ethyl 2-benzylideneamino-2-ethylhexanoate

Sodium hydride (32.5g, 60% dispersion in oil) and N,N-dimethylformamide (DMF) (700ml) were stirred under nitrogen at room temperature and a solution of the product from step (b) (178.1g) in DMF was added dropwise. After 2 hours stirring at room temperature, a solution of butyl iodide (149.5g) in DMF was added dropwise and the reaction left stirring for a further 2 hours. The reaction was poured into an ice cold mixture of water (560ml), diethyl ether (300ml) and ammonium chloride (120g). The resulting organic layer was dried over potassium carbonate then concentrated to give the desired product as a brown oil (220g).

(d) (±)-Ethyl 2-amino-2-ethylhexanoate

The product from step (c) (233.0g) was partitioned between petroleum ether and 10% w/w hydrochloric acid (421ml) and stirred at room temperature for 2 hours. The aqueous layer was extracted twice with petroleum ether and then chilled with ethyl acetate in an ice-salt bath. Sodium hydroxide pellets were added to the mixture until the aqueous layer was at pH 10. The latter was extracted twice with ethyl acetate and the combined ethyl acetate layers were dried over potassium carbonate, then concentrated and vacuum distilled to give the desired product as a colourless oil. ¹H NMR consistent with the proposed structure.

(e) (R)-2-Amino-2-ethylhexanoic acid

A suspension of pig liver esterase (0.1g, Sigma-Aldrich-Fluka) in water was added to an aqueous solution of the product from step (d) (100g). When addition was complete, the pH of the mixture was adjusted to 9.7 using 1N aqueous NaOH and maintained at this value by the addition of further 1N NaOH. After the addition of a predetermined amount of 1N aqueous NaOH (85g over 10 hours), the mixture was washed with diethyl ether to remove unreacted (S)-ethyl 2-amino-2-ethylhexanoate. The remaining aqueous phase was evaporated in vacuo to give a white solid comprising the titled compound and its sodium salt.

21

(f) (R)-2-Amino-2-ethylhexan-1-ol

The product (20g) from step (e) was added to a 1M solution of lithium aluminium hydride (1.5 molar equivalents) in THF and the mixture was refluxed for 3 hours, then stirred for 16 hours at room temperature. The mixture was cooled to about 0°C, then quenched with water and 1N aqueous NaOH. The resulting solid was broken up with additional water and the suspension was heated at 50°C for 5 minutes, then cooled to room temperature. Diethyl ether (100ml) was added, the mixture was stirred and filtered. The diethyl ether layer was separated, dried and concentrated in vacuo to give the desired product as an oil (82% yield). ¹H NMR consistent with the proposed structure.

(g) (R)-2-Amino-2-ethylhexyl hydrogen sulfate

The product (20.0g) from step (f) was dissolved in dichloromethane (170 ml) and treated with chlorosulfonic acid (26.8 g). The reaction mixture was stirred at room temperature for 17 hours. A major part of the solvent was removed by distillation and the resulting slurry was diluted with acetone, filtered and dried to get a white solid. ¹H NMR consistent with the proposed structure.

(h) 2-Hydroxy-4,5-dimethoxybenzaldehyde

A 1.0M solution of boron trichloride (210 ml, Aldrich) in dichloromethane was added to benzoyl chloride (30.1g, Aldrich) in benzene (350 ml). Next, 3,4-dimethoxyphenol (30.0g, Aldrich) in benzene (130 ml) was added and the reaction mixture was stirred at room temperature for 2 1/2 hours. 50% NaOH (55 ml) was then added and the mixture was stirred for 15 minutes. The organic layers were separated, dried and concentrated in vacuo. The resulting residue was triturated with 1N NaOH for 40 minutes then filtered. The aqueous basic filtrate was acidified with conc. HCl to give the title product as a yellow solid (25.9g), mp 104-105°C. ¹H NMR was consistent with the proposed structure.

(i) O-(2-Benzoyl-4,5-dimethoxyphenyl)-N,N-diethylthiocarbamate

Triethylamine (106.3g, Aldrich), 4-dimethylaminopyridine (6.5g, Aldrich) and diethylthiocarbamoyl (86.4g) was added to the product (130.4g) from step(h) to 1 L of dioxane. The reaction mixture was stirred at reflux for 22 hours, cooled to room temperature, then filtered. The filtrate was concentrated in vacuo and 1N HCl (600 ml) was added followed by diethyl ether (500 ml). The mixture was allowed to stand for 45

22

minutes, then filtered. The solids were washed thoroughly with diethyl ether and dried in a vacuum oven to afford the title product as a yellow solid (120.5g), mp 94-95°C. ¹H NMR was consistent with the proposed structure.

(j) S-(2-Benzoyl-4,5-dimethoxyphenyl)-N,N-diethylthiocarbamate

A slurry of the product (60.4g) from step (i) in tetradecane (250 ml) was heated to an internal temperature of 250°C and kept there for a period of 25 minutes. The reaction mixture was cooled with an ice bath. The solvent was decanted and the residue was triturated with diethyl ether (100 ml) to give the title product (43.9g) as a beige solid, mp 114-116°C. ¹H NMR was consistent with the proposed structure.

(k) 2-Citricaplo-4,5-dimethoxybenzophenone

Potassium hydroxide pellets (58.6g) was slowly added to a solution of the product (85.0g) from step (j) dissolved in 1L of methanol/THF (1:1). After refluxing for 3 hours, the reaction mixture was cooled to room temperature and concentrated in vacuo. The resulting residue was triturated with 1N HCl then extracted with EtOAc. The organic layer was separated, washed with 2x250 ml of 1N HCl then washed with 3x400 ml of 1N NaOH. The aqueous basic layers were combined and acidified with conc. HCl to afford the title product (54.8g) as a gold solid. ¹H NMR was consistent with the proposed structure.

(l) (R)-2-(2-Amino-2-ethylhexylthio)-4,5-dimethoxybenzophenone

The product (48.8g) from step (g) was dissolved in water (250 ml) and to this solution the product (54.2g) from step (k) in butyl acetate (300 ml) was added. The reaction mixture was stirred and heated to an internal temperature of 95°C and NaOH (18.9g) in water (250 ml) was added dropwise. After complete addition, the reaction was stirred an additional 25 minutes at 95°C, then cooled to room temperature. The organic layer was separated, dried and concentrated to give the title product (78.5g) as an orange-brown oil. Treatment of the free base with ethereal HCl afforded the hydrochloride salt as a light yellow solid, mp 75-78°C. ¹H NMR consistent with the proposed structure.

(m) (3R)-3-Butyl-3-ethyl-2,3-dihydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine

The product (78.0g) from step (l) was dissolved in 2,6-lutidine (400ml), added p-toluenesulfonic acid (0.70g) and the reaction mixture was refluxed using a Dean Stark trap.

The reaction was refluxed for a period of 22 hours during which time solvent was removed from the apparatus and then replaced with fresh solvent. The reaction mixture was concentrated in vacuo and the residue was treated with 5% NaHCO₃(300ml) and EtOAc(300ml). The EtOAc layer was separated, washed with brine, dried and concentrated in vacuo to give a dark red oil. Chromatography on silica gel, using hexane : EtOAc(4:1) as eluant, afforded the desired product (64.1g) as a light brown oil. ¹H NMR consistent for the proposed structure.

(n) (1R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine

A 1M solution of diborane in THF (200ml) was added to a solution of the product (64.0g) from step (m) in THF (350ml). The reaction mixture was stirred at room temperature for a period of 17 hours, then 6N HCl (150ml) was added and the solution was concentrated in vacuo to remove THF. The aqueous residue was basified with 50% NaOH and extracted with EtOAc. The EtOAc layer was separated, dried and concentrated in vacuo to afford an oil which was chromatographed on silica gel, using hexane : EtOAc (85:15) as eluant, to give the title product (25.5g) as a beige solid, mp 64-66°C. ¹H NMR consistent for proposed structure.

(o) (1R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide

A solution of the product (25.5g) from step (n) in trifluoroacetic acid (125ml) was added to 30% H₂O₂ (18.8g) in trifluoroacetic acid (100ml). The reaction mixture was stirred at room temperature for 17 hours, then poured into water (800ml) followed by the addition of 50% NaOH until the mixture reached a pH of 10. The reaction mixture was layered with EtOAc and stirred for 1 hour. The organic layer was separated, dried and concentrated in vacuo to afford solids which were recrystallized from EtOH to afford the title product (18.5g) as a white solid, mp 148-149°C.

Analysis: Calcd : C 66.16; H 7.48; N 3.35; S 7.68
Found: C 66.01; H 7.56; N 3.31; S 7.74

¹H NMR (DMSO-d₆) δ: 0.74-0.86(6H, m); 1.07-1.39 (4H, m); 1.39-2.20 (4H, m); 3.33 (2H, q); 3.44 (3H, s); 3.83 (3H, s); 5.92 (1H, d); 6.11 (1H, s); 7.33-7.48 (6H, m).

Synthetic Example 2

Preparation of (1R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-4-ol 1,1-dioxide

Oxone (146.7g Aldrich) in water (550 ml) was added to a solution of the product (18.4g) from Synthetic Example 1(n) in MeOH (500ml). The reaction mixture was stirred at room temperature for 17 hours, then cautiously basified with 50% NaOH. This heterogeneous mixture was layered with EtOAc and stirred for 1 hour. The organic layer was separated, dried and concentrated in vacuo to get a pink solid. Chromatography on silica gel, using hexane: EtOAc (65:35) as eluant, afforded the title product (6.7g) as a white solid, mp 174-175°C.

Analysis: Calcd: C 63.72; H 7.21; N 3.23; S 7.39
Found: C 63.81; H 7.22; N 3.19; S 7.47

¹H NMR(DMSO-d₆) δ: 0.77-0.90 (6H, m); 1.10-2.17 (8H, m); 3.27-3.45 (5H, m); 3.84 (3H, s); 6.14 (1H, s); 6.38 (1H, s); 7.30-7.53 (5H, m); 7.97 (1H, s).

Synthetic Example 3

Preparation of (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide

(a) (±)-2-Amino-2-ethylhexan-1-ol

Lithium aluminum hydride (22.2g) was added to anhydrous diethyl ether (450ml) under nitrogen. The product (129.0g) from Synthetic Example 1(d) was diluted with diethyl ether (40ml) and added dropwise. The reaction was refluxed for 1 hour then cooled to room temperature. 1M sodium hydroxide (23ml) was added dropwise followed by deionised water. The resulting suspension was filtered and the filtrate concentrated to give the desired product as a colorless oil (87.9g). ¹H NMR consistent with the proposed structure.

(b) (±)-2-Butyl-2-ethylaziridine

Acetonitrile (150ml) and the product (20.0g) from step (a) were mixed under nitrogen, cooled to 2-3°C and chlorosulphonic acid (16.0g) Aldrich) was added dropwise keeping the temperature below 10°C. The coolant was removed and the slurry left to stir for 80

25

minutes at room temperature. The reaction was concentrated in vacuo and co-distilled with water (50ml). 50% Aqueous sodium hydroxide (55.2g) and water (50ml) were added and the mixture was distilled at atmospheric pressure. The organic layer was collected from the distillate and dried with solid potassium hydroxide to give the desired product (12.8g). ¹H NMR consistent with proposed structure.

(c) (±)-3-Buryl-3-ethyl-2,3-dihydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine

The product (14.7g) from Synthetic Example 1(k), in 2,6-lutidine (50ml), was added to a solution of the product (6.5g) from step (b) in 2,6-lutidine (200ml). The reaction mixture was stirred for 1 hour, conc. HCl (4.4ml) was added and then refluxed with a Dean-Stark trap for 17 hours. The reaction mixture was concentrated in vacuo and the residue was partitioned between 5% NaHCO₃ and EtOAc. The organic layer was separated, dried and concentrated to get an oil which was chromatographed on silica gel, using hexane: EtOAc (7:3) as eluant to afford the desired product (12.0g) as an oil. ¹H NMR consistent with proposed structure.

d) (±)-[Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide]

The title compound was prepared following the procedures of Synthetic Example 1 (n)-(o) using the product from step (c) to give a white solid, mp 146-147°C.

Analysis (0.50 H₂O): Calcd: C 64.54; H 7.35; N 3.24; S 7.40

Found: C 64.76; H 7.56; N 3.28; S 7.52

¹H NMR (DMSO-d₆): δ: 0.74-0.86 (6H, m); 1.07-1.39 (4H, m); 1.40-2.20 (4H, m); 3.33 (2H, q); 3.44 (3H, s); 3.83 (3H, s); 5.92 (1H, d); 6.11 (1H, s); 7.30-7.48 (6H, m)

Synthetic Example 4

Preparation of (±)-[Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-4-ol 1,1-dioxide]

Oxone (7.3g, Aldrich) in water (100ml) was added to a solution of the product (1.7g) from Synthetic Example 3(d) in MeOH (100ml). The reaction mixture was stirred at room temperature for a period of 17 hours and water and EtOAc were added. After stirring for 1 hour, the organic layer was separated, dried and concentrated to give a foam.

26

Chromatography on silica gel, using hexane: EtOAc (4:1) as eluant, gave the desired product (1.2g) as a white solid, mp 172-174°C.

Analysis: Calcd: C 63.72; H 7.21; N 3.23; S 7.39

Found: C 63.79; H 7.26; N 3.18; S 7.47

¹H NMR (DMSO-d₆): δ: 0.78-0.90 (6H, m); 1.14-2.14 (8H, m); 3.27-3.41 (5H, m); 3.84 (3H, s); 6.13 (1H, s); 6.37 (1H, s); 7.34-7.53 (5H, m); 7.96 (1H, s).

Synthetic Example 5

Preparation of (3R,5R)-7-Bromo-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide

(a) O-(2-benzoyl-5-methoxyphenyl)dimethylthiocarbamate

Sodium hydride (8.8g, Aldrich) was added slowly to a solution of 2-hydroxy-4-methoxybenzophenone (50.0g, Aldrich) in 300 ml of dimethylformamide. Hexamethylphosphoramide (43.0g) was then added dropwise and stirred at room temperature for 2 hours. Dimethylthiocarbonyl chloride (37.0g, Aldrich) was added and stirred overnight at 50°C. The reaction mixture was poured into deionized water (300ml) and extracted with a petroleum ether/chloroform (1:4) mixture. The organic layer was washed with 10% sodium hydroxide, brine and concentrated to give the title product as a yellow solid (40.0g). ¹H NMR was consistent with proposed structure.

(b) S-(2-Benzoyl-5-methoxyphenyl)dimethylthiocarbamate

The product (97.4g) from step (a) was suspended in tetradecane (500ml) and heated to an internal temperature of 255°C for 30 minutes. After cooling to room temperature, the reaction mixture was chromatographed on silica gel using hexane, then hexane/ethyl acetate (7:3) as eluants to afford the title product (65.0g) as a tan solid, mp 95-97°C. ¹H NMR was consistent with proposed structure.

(c) 2-Mercapto-4-methoxybenzophenone

Potassium hydroxide pellets (20.0g) were slowly added to a solution of the product (28.0g) from step (b) dissolved in 800 ml methanol/tert-butylhydrofuran (1:1). After refluxing for 4 hours, the reaction was cooled to room temperature, methylene chloride was added and the solution was extracted with 5% hydrochloric acid. The organic layer was dried and

27

concentrated. Chromatography on silica gel using hexanes/ethyl acetate (99:1) as the eluant afforded the title product (17.1g) as an orange oil. ¹H NMR consistent with proposed structure.

(d) (R)-2-(2-Amino-2-ethylhexylthio)-4-methoxybenzophenone

This compound was prepared following the procedure of Synthetic Example 1(i), using the product (46.4g) from step (c) and the product (44.6g) from Synthetic Example 1(g). Concentration of the organic layer afforded the title product (66.5g) as a red oil. ¹H NMR consistent with proposed structure.

(e) (2R,5R)-2-Butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine

This compound was prepared following the procedure of Synthetic Example 1(m), using the product (66.5g) from step (d). Chromatography on silica gel, using hexane: EtOAc (9:1) as eluant, afforded the title compound (54.5g) as a yellow oil. ¹H NMR consistent with the proposed structure.

(f) (2R,5R)-2-Butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine

This compound was prepared following the procedure of Synthetic Example 1(n), using the product (54.4g) from step (e). Chromatography on silica gel, using hexane:EtOAc (9:1) as eluant, gave the title product (22.8g) as an orange oil. ¹H NMR consistent with the proposed structure.

(g) (2R,5R)-2-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine

Bromine (18.6g) was added to a solution of the product (10.4g) from step (f) dissolved in glacial acetic acid (150ml). The reaction mixture was stirred at room temperature for 2 hours. Acetic acid was removed in vacuo, added another 100ml and concentrated in vacuo. The resulting residue was dissolved in EtOAc and washed with sodium metabisulfite and 1N NaOH. The organic layer was separated, dried and concentrated in vacuo to give a brown oil which was then converted to the hydrochloride salt with ethereal HCl. This solid was filtered, washed with ether and then treated with 1N NaOH and EtOAc

28

to get the title product (8.9g) as an orange oil. ¹H NMR consistent with the proposed structure.

(h) (2R,5R)-2-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide

This compound was prepared following the procedure of Synthetic Example 1(o), using the product (8.2g) from step (g). Chromatography on silica gel, using hexane: EtOAc (4:1) as eluant, afforded a foam which upon trituration in ether gave the title product (5.0g) as a white solid, mp 132-134°C.

Analysis: Calcd: C 56.65; H 6.05; N 3.00; Br 17.13; S 6.87

Found: C 56.71; H 6.01; N 2.94; Br 17.07; S 6.95

¹H NMR (DMSO-d₆): δ: 0.64-0.81 (6H, m); 0.97-1.19 (4H, m); 1.22-1.50 (2H, m); 1.69-1.78 (1H, m); 1.98-2.06 (1H, m); 2.67 (1H, d); 3.39 (2H, q); 3.92 (3H, s); 5.88 (1H, d); 6.63 (1H, s); 7.29-7.43 (5H, m); 7.55 (1H, s)

Synthetic Example 6

Preparation of (2R,5R)-2-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide

Meta-chloroperoxybenzoic acid (57-86%, 0.90g, Aldrich) in CH₂Cl₂ (50ml) was added to a solution of the product (2.4g) from Synthetic Example 5(h) in CH₂Cl₂ (50ml). The reaction was stirred at room temperature for 1 hour, then 5% NaHCO₃ (100ml) was added and the mixture stirred for 30 minutes. The organic layer was separated, dried and concentrated in vacuo to give a foam. Chromatography on silica gel, using hexane: EtOAc (9:1) as eluant gave a foam which upon trituration with ether afforded the title product (1.3g) as a white solid, mp 202-204°C.

Analysis: Calcd: C 54.77; H 5.85; N 2.90; Br 16.56; S 6.65

Found: C 54.92; H 5.90; N 2.85; Br 16.65; S 6.75

¹H NMR (DMSO-d₆): δ: 0.75-0.86 (6H, m); 1.05-1.41 (5H, m); 1.43-1.64 (1H, m); 1.66-1.79 (1H, m); 1.83-2.49 (1H, m); 3.46 (2H, s); 3.93 (3H, s); 6.33 (1H, s); 6.67 (1H, s); 7.30-7.50 (6H, m); 8.07 (1H, s)

29

Synthetic Example 7

Preparation of (3R,5R)-3-Benzyloxy-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-7,8-diol-1,1-dioxide

The product (5.0g) from Synthetic Example 1(c) was dissolved in glacial acetic acid (36 ml) and 48% HBr (36ml) and allowed to stir at reflux for 2 hours. The reaction mixture was poured into an ice water mixture then basified with 50% NaOH to a pH of 7. The reaction mixture was filtered to get a solid which was chromatographed on silica gel, using hexane: EtOAc (3:2) as eluant, to give the title product (1.6g) as a white solid, mp 117-118°C.

Analysis (0.30 H₂O): Calcd: C 63.87; H 7.04; N 3.55; S 8.12

Found: C 63.86; H 7.09; N 3.51; S 8.18

¹H NMR (DMSO-d₆): δ: 0.76 (3H, t); 0.81 (3H, t); 1.08-2.41 (8H, m); 3.24 (2H, q); 5.83 (1H, d); 6.03 (1H, s); 7.31-7.42 (6H, m); 9.60 (3H, bs)

Synthetic Example 8

Preparation of (3R,5R)-3-Benzyloxy-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-ol-1,1-dioxide

Chromatography of the reaction from Synthetic Example 7 produced mixtures which were combined and rechromatographed using toluene and toluene: EtOAc (95:5) as the eluants, to afford the title product (0.29g) as a white solid, mp 155-156°C.

Analysis: Calcd: C 65.48; H 7.24; N 3.47; S 7.95

Found: C 65.58; H 7.28; N 3.43; S 8.03

¹H NMR (DMSO-d₆): δ: 0.76 (3H, t); 0.81 (3H, t); 1.18-2.04 (8H, m); 3.28 (2H, q); 3.82 (3H, s); 5.85 (1H, d); 6.09 (1H, s); 7.31-7.45 (6H, m); 9.43 (1H, s)

Synthetic Example 9

Preparation of (3R,5R)-3-Benzyloxy-2,3,4,5-tetrahydro-7-methoxy-5-phenyl-1,4-benzothiazepine-8-ol-1,1-dioxide

Chromatography of the reaction mixtures from Synthetic Example 7 produced the title compounds of Synthetic examples 7 and 8. One other product was also isolated during the chromatography used in Synthetic Example 8. The title product (0.35g) was isolated as a white solid, mp 165-166°C.

30

Analysis: Calcd: C 65.48; H 7.24; N 3.47; S 7.95

Found: C 65.32; H 7.28; N 3.49; S 8.00

¹H NMR (DMSO-d₆): δ: 0.77 (3H, t); 0.81 (3H, t); 1.11-2.08 (8H, m); 3.29 (2H, q); 3.44 (3H, s); 5.86 (1H, d); 6.06 (1H, s); 7.32-7.43 (6H, m); 9.73 (1H, s)

Synthetic Example 10

Preparation of (±)-trans-3-benzyloxy-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide

(a) O-(2-benzyloxy-5-methoxyphenyl)dimethylthiocarbamate

Sodium hydride (8.8g, Aldrich) was added slowly to a solution of 2-hydroxy-4-methoxybenzophenone (50.0g Aldrich) in 300 ml of dimethylformamide. Hexamethylphosphoramide (43.0g) was then added dropwise and stirred at room temperature for 2 hours. Dimethylthiocarbamoyl chloride (37.0g Aldrich) was added and stirred overnight at 50°C. The reaction mixture was poured into deionized water (300ml) and extracted with a petroleum ether/chloroform (1:4) mixture. The organic layer was washed with 10% sodium hydroxide, brine and concentrated to give the title product as a yellow solid (40.0g). ¹H NMR was consistent with proposed structure.

(b) N-(2-benzyloxy-5-methoxyphenyl)dimethylthiocarbamate

The product (97.4g) from step (a) was suspended in tetradecane (500ml) and heated to an internal temperature of 255°C for 30 minutes. After cooling to room temperature, the reaction mixture was chromatographed on silica gel using hexane, then hexanes/ethyl acetate (7:3) as eluants to afford the title product (65.0g) as a tan solid, mp 95-97°C. ¹H NMR was consistent with proposed structure.

(c) 2-Mercapto-4-methoxybenzophenone

Potassium hydroxide pellets (20.0g) were slowly added to a solution of the product (28.0g) from step (b) dissolved in 800 ml methanol/tetrahydrofuran (1:1). After refluxing for 4 hours, the reaction was cooled to room temperature, methylene chloride was added and the solution was extracted with 5% hydrochloric acid. The organic layer was dried and concentrated. Chromatography on silica gel using hexanes/ethyl acetate (99:1) as the eluant afforded the title product as an orange oil (17.1g). ¹H NMR consistent with proposed structure.

(d) Ethyl 2-aminobutyrate hydrochloride

A slurry of 2-aminobutyric acid (100g Aldrich) in absolute ethanol (300ml) was stirred under nitrogen at 0°C and thionyl chloride (120.8g) was added dropwise. The reaction was stirred overnight at 0°C and then gradually warmed to room temperature. The resulting white slurry was heated under reflux for 3 hours, left to cool for 10 minutes, then poured into chilled diethyl ether (600ml) with hand stirring. The suspension was filtered and the solid product dried to give the desired product (150g) as a white solid. ¹H NMR consistent with proposed structure.

(e) Ethyl 2-benzylidenemino-2-ethylhexanoate

A solution of the product from step (d) (149.6g), magnesium sulphate (74.3g), and triethylamine (246ml) in dichloromethane (1500ml) was stirred at room temperature under nitrogen and benzaldehyde (94.9g, Aldrich) was added dropwise. The mixture was stirred at room temperature for 3 hours then filtered. The filtrate was concentrated, triturated in diethyl ether, filtered and concentrated to yield the desired product as a yellow oil (174g). ¹H NMR consistent with the proposed structure.

(f) (±)-Ethyl 2-benzylidenemino-2-ethylhexanoate

Sodium hydride (32.5g, 60% dispersion in oil) and N,N-dimethylformamide (DMF) (700ml) were stirred under nitrogen at room temperature and a solution of the product from step (e) (178.1g) in DMF was added dropwise. After 2 hours stirring at room temperature, a solution of butyl iodide (149.5g) in DMF was added dropwise and the reaction left stirring for a further 2 hours. The reaction was poured into an ice cold mixture of water (560ml), diethyl ether (300ml) and ammonium chloride (120g). The resulting organic layer was dried over potassium carbonate then concentrated to give the desired product as a brown oil (220g).

(g) (±)-Ethyl 2-amino-2-ethylhexanoate

The product from step (f) (233.0g) was partitioned between petroleum ether and 10% w/w hydrochloric acid (421ml) and stirred at room temperature for 2 hours. The aqueous layer was extracted twice with petroleum ether and then chilled with ethyl acetate in an ice-salt bath. Sodium hydroxide pellets were added to the mixture until the aqueous layer was at

pH 10. The latter was extracted twice with ethyl acetate and the combined ethyl acetate layers were dried over potassium carbonate, then concentrated and vacuum distilled to give the desired product as a colourless oil. ¹H NMR consistent with the proposed structure.

(h) (±)-2-Amino-2-ethylhexan-1-ol

Lithium aluminium hydride (22.2g) was added to anhydrous diethyl ether (450ml) under nitrogen. The product from step (g) (129.0g) was diluted with diethyl ether (40ml) and added dropwise. The reaction was refluxed for 1 hour then cooled to room temperature. 1M sodium hydroxide (23ml) was added dropwise followed by deionised water. The resulting suspension was filtered and the filtrate concentrated to give the desired product as a colourless oil (87.9g). ¹H NMR consistent with the proposed structure.

(i) (±)-2-Buryl-2-ethylaziridine

Acetonitrile (150ml) and the product from step (h) (20.0g) were mixed under nitrogen, cooled to 2-3°C and chlorosulphonic acid (16.0g, Aldrich) was added dropwise keeping the temperature below 10°C. The coolant was removed and the slurry left to stir for 80 minutes at room temperature. The reaction was concentrated in vacuo and co-distilled with water (50ml). 50% Aqueous sodium hydroxide (55.2g) and water (50ml) were added and the mixture was distilled at atmospheric pressure. The organic layer was collected from the distillate and dried with solid potassium hydroxide to give the desired product (12.8g). ¹H NMR consistent with proposed structure.

(j) (±)-3-Buryl-3-ethyl-8-methoxy-5-phenyl-2,3-dihydrobenzothiazepine

The product (55.2g) from step (i), in 2,6-lutidine (100ml), was added to a solution of the product (118.5g) from step (e) in 2,6-lutidine (400ml). The reaction mixture was stirred for 1 hour and p-toluenesulphonic acid (9.0g) was added and then refluxed with a Dean Stark trap for 17 hours. The reaction mixture was concentrated in vacuo and the residue was partitioned between 5% NaHCO₃ and EtOAc. The organic layer was separated, dried and concentrated to get an oil which was chromatographed on silica gel, using hexane: EtOAc (85:15) as the eluant, to afford the title product (124.3g) as an orange oil. ¹H NMR consistent with the desired structure.

33

(k) (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine

A 1M solution of diborane (40ml) in THF was added to a solution of the product (12.3g) from step (j) in THF (150ml). The reaction mixture was stirred at room temperature for 17 hours, then 6N HCl (50ml) was added and the solution was concentrated in vacuo. The residue was basified with 50% NaOH and extracted with EtOAc. The EtOAc layer was separated, dried and concentrated in vacuo to give an oil which was chromatographed on silica gel, using hexanes then toluene as the eluants to afford the desired product (4.9g) as a yellow oil. ¹H NMR consistent with the desired structure.

(l) (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide

A solution of the product (4.9g) from step (k) in trifluoroacetic acid (50ml) was added to 30% H₂O₂ in trifluoroacetic acid (50ml). The reaction mixture was stirred at room temperature for 17 hours, then poured into deionized water (200ml) followed by the addition of NaOH pellets until a pH of 14 was obtained. The reaction mixture was warmed, at 45°C, for 3 hours then extracted with dichloromethane. The organic layer was separated, dried and concentrated to give an oil which was chromatographed on silica gel, using hexane : EtOAc (9:1) as the eluant, to give the title product as a white solid, mp 123-125°C.

Analysis: Calcd: C 68.18; H 7.54; N 3.61; S 8.27

Found: C 68.19; H 7.49; N 3.55; S 8.35

¹H NMR (DMSO-d₆): δ: 0.73-0.85 (6H, m, CH₃); 1.07-1.47 (4H, m, CH₂); 1.48-2.20 (4H, m, CH₂); 2.48-2.53 (1H, d, NH); 3.51 (2H, q, CH₂SO₂); 3.84 (3H, s, OMe); 5.90 (1H, d, CHPh); 6.50 (1H, d, ArH); 7.09-7.20 (1H, m, ArH); 7.32-7.48 (6H, m, ArH)

Synthetic Example 11

Preparation of (+)-trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-ol 1,1-dioxide

This compound was prepared following the procedure of Synthetic Example 7, using the product (4.8g) from Synthetic Example 10(1). Chromatography on silica gel, using hexane: EtOAc (4:1) as the eluant, gave the title product (1.8g) as a white solid, mp 130-132°C.

34

Analysis: Calcd: C 67.53; H 7.28; N 3.75; S 8.58
Found: C 67.26; H 7.21; N 3.76; S 8.65

¹H NMR (DMSO-d₆): δ: 0.70-0.86 (6H, m); 0.96-1.23 (4H, m); 1.25-1.49 (1H, m); 1.66-1.75 (1H, m); 1.98-2.07 (1H, m); 2.40 (1H, d); 3.33 (2H, q); 5.82 (1H, d); 6.35 (1H, d); 6.77-6.80 (1H, m); 7.24-7.38 (6H, m); 10.0 (1H, s)

Synthetic Example 12

Preparation of (+)-trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-4,8-diol 1,1-dioxide

The product (1.0g) from Synthetic Example 11 was dissolved in methylene chloride (100ml), cooled to 0°C and m-chloroperoxybenzoic acid (0.55g, 57-86%, Aldrich) was added. The reaction mixture was stirred at ice bath temperatures for 5 hours then 5% NaHCO₃ was added to neutralize excess acid. The organic layer was separated, dried and concentrated in vacuo. The resulting residue was chromatographed on silica gel, using hexane: EtOAc as the eluant, to afford the title product (0.68g) as a pale yellow solid, mp 213-214°C.

Analysis: Calcd: C 64.76; H 6.99; N 3.60; S 8.23

Found: C 64.86; H 7.03; N 3.63; S 8.31

¹H NMR (DMSO-d₆): δ: 0.77-0.89 (6H, m); 1.09-1.64 (6H, m); 1.68-2.03 (2H, m); 3.36 (2H, q); 6.30 (1H, s); 6.44 (1H, d); 6.82-6.87 (1H, m); 7.27-7.49 (6H, m); 7.89 (1H, s); 10.0 (1H, s)

Synthetic Example 13

Preparation of (+)-trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methyl-5-phenyl-1,4-benzothiazepine 1,1-dioxide

(a) 3-Methylphenyl benzoate

A solution of benzoyl chloride (32.5g, Aldrich) in ether (200ml) was added dropwise to a stirred solution of m-cresol (25.0g, Aldrich) and triethylamine (27.2g, Aldrich) in ether (500ml). The reaction mixture was stirred at room temperature for 1 hour then filtered. The etheral filtrate was washed with saturated NaHCO₃ and water then dried over Na₂SO₄. The ether layer was separated, dried and concentrated in vacuo to give the

desired product (104.0g) as a white solid, mp 45-47°C. ¹H NMR consistent with the desired structure.

(b) 2-Hydroxy-4-methylbenzophenone

The product (48g) from step (a) was melted (at 70°C) and aluminum chloride (30.2g) was added in portions. The reaction mixture was heated to 200°C for 5 minutes, then cooled to room temperature. The resulting solid was ground to a powder and slowly added to a mixture of conc. HCl (800ml) and ice. This mixture was extracted with ether and the ether was washed with water. The ether layer was separated, dried and concentrated. The resulting residue was chromatographed on silica gel, using toluene as the eluant, to afford the title product (39g) as a yellow oil. ¹H NMR consistent with the desired structure.

(c) (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methyl-5-phenyl-1,4-benzothiazepine 1,1-dioxide

The product from step (b) was converted to the title product following the procedures used in steps (a) to (f) of Synthetic Example 10. The title product was isolated as a white solid, mp 121-122°C

Analysis: C 71.12; H 7.87; N 3.77; S 8.63
Found: C 71.23; H 7.94; N 3.67; S 8.74

¹H NMR (DMSO-d₆): δ: 0.77-0.82 (6H, m); 1.16-2.07 (8H, m); 2.36 (3H, s); 3.37 (2H, q); 5.92 (1H, d); 6.47 (1H, d); 7.27-7.39 (6H, m); 7.79 (1H, s)

Synthetic Example 14

Preparation of (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide

(a) (±)-2-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (16.9g) was added directly to a benzene solution (300 ml) of the product (as the racemate) (30.2g) from Synthetic Example 7 (g). The reaction mixture was stirred at reflux for 3 hours, cooled to room temperature. 1N NaOH (200 ml) was added, stirred for 30 minutes, then the organic layer was separated, washed with brine and 1N NaOH. The benzene layer was separated, dried and concentrated

to get an oil which was solubilized in hexane, filtered and concentrated to get the title product (25.8g) as a red oil. ¹H NMR consistent with the desired structure.

(b) (±)-3-Butyl-7-carbaldehyde-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine

A solution of 1.6M n-butyl lithium (49.0 ml) was to an ice-cooled solution of the product (25.8g) from step (a) in hexane (500 ml). The reaction mixture was stirred for 25 minutes and 4-formylmorpholine (9.0g). The ice bath was removed and the reaction was stirred at room temperature for 2 1/2 hours. The reaction was quenched with a saturated solution (250 ml) of NH₄Cl and stirred for 60 minutes. The organic layer was separated, dried and concentrated to get 26.9g of a red oil. Chromatography on silica gel, using hexane:EtOAc (85:15) as eluant afforded the title product (13.9g) as an orange oil. ¹H NMR consistent with the desired structure.

(c) (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde

Ethylene glycol (9.3g) and pyridinium p-toluenesulfonate (1.3g) were added to a benzene solution (250 ml) of the product (19.0g) from step (b) and this mixture was refluxed in a Dean Stark trap for 17 hours. The reaction mixture was cooled to room temperature and treated with aqueous NaHCO₃ (150 ml) for 15 minutes. The organic layer was separated, dried and concentrated to get a thick yellow-orange oil (19.7g). ¹H nmr was consistent for the dioxolane derivative. This oil was then treated with B₂H₆, following the procedure cited in Synthetic Example 1 (n), to give the title product (3.5g) as an orange oil. ¹H NMR consistent with the desired structure.

(d) (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide

The product (3.5g) from step (c) was dissolved in t-butanol/THF (1:4, 60 ml) and N-methylmorpholine N-oxide (3.4g) was added, followed by 2.5 wt.% of OsO₄ in 2-methyl-2-propanol (5.0 ml). The reaction mixture was stirred at room temperature for 17 hours then diluted with EtOAc (250 ml). The organic layer was separated, washed with 1N NaOH (2x150 ml) and brine. The organic layer was separated, dried and concentrated to give an oil which upon trituration in diethyl ether afforded the title product (3.10g) as a white solid, mp 127-128°C.

37

Analysis: Calcd: C 66.48; H 7.03; N 3.37; S 7.72
Found: C 66.26; H 7.04; N 3.30; S 7.82

¹H NMR(DMSO-d₆): δ: 0.73-0.86(6H, m); 1.07-2.05(8H, m); 2.65(1H, d); 3.50(2H, q); 4.03(3H, s); 5.91(1H, s); 6.92(1H, s); 7.33-7.48(5H, m); 7.74(1H, s); 10.28(1H, s)

Synthetic Example 15

Preparation of (+)-Trans-2-(3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepin-7-yl)methoxy) ethanol S,S-dioxide

Chromatography of the reaction from Synthetic Example 14 (c) afforded the corresponding sulfide compound(2.3g) of the title product as an oil. ¹H NMR was consistent for the desired structure. This oil was then treated, according to the procedure shown in Synthetic Example 1 (c), to yield the title product (0.65g) as a white solid, mp 83-85° C.

Analysis: Calcd: C 65.05; H 7.64; N 3.03; S 6.95
Found: C 64.82; H 7.72; N 2.99; S 6.91

¹H NMR(DMSO-d₆): δ: 0.74-0.86(6H, m); 1.07-2.14(8H, m); 2.52(1H, d); 3.35(4H, m); 3.41(2H, q); 3.87(3H, s); 4.39(2H, s); 4.54(1H, t); 5.91(1H, d); 6.64(1H, s); 7.29-7.45(5H, m); 7.51(1H, s)

Synthetic Example 16

Preparation of (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-hydroxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide

The product (2.0g) from Synthetic Example 14 (d) was added to glacial acetic acid (20 ml) and 48% HBr (20 ml) and heated to 150° C for 24 hours. The reaction mixture was concentrated in vacuo, partitioned between diethyl ether and 5% NaHCO₃. The organic layer was separated, dried and concentrated to give the title product (0.85g) as a tan solid, mp 158-159° C.

Analysis: Calcd: C 65.81; H 6.78; N 3.49; S 7.99
Found: C 65.63; H 7.04; N 3.32; S 7.74

38

¹H NMR(DMSO-d₆): δ: 0.72-0.85(6H, m); 1.07-2.05(8H, m); 2.58(1H, d); 3.46(2H, q); 5.85(1H, d); 6.83(1H, s); 7.34-7.47(5H, m); 7.70(1H, s); 10.25(1H, s); 11.33(1H, broad s)

Synthetic Example 17

Preparation of (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-thiol 1,1-dioxide

The product from Synthetic Example 11 was treated according to the procedures shown in Synthetic Example 1 (i)-(k) to give the title product as a white solid, mp 108-110° C.

Analysis: Calcd: C 64.75; H 6.99; N 3.60; S 16.46
Found: C 64.83; H 7.03; N 3.56; S 16.54

¹H NMR(DMSO-d₆): δ: 0.70-0.81(6H, m); 1.05-2.06(8H, m); 2.34(1H, d); 3.37(2H, q); 5.85(1H, d); 6.06(1H, broad s); 6.40(1H, d); 7.26-7.40(6H, m); 7.90(1H, s)

Synthetic Example 18

Preparation of (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-sulfonic acid 1,1-dioxide

The product(5.3g) from Synthetic Example 17 was dissolved in DMSO (13 ml). Water (0.3 ml) and 48% HBr (0.2 ml) were then added. The reaction mixture was heated to 120° C, allowing for distillate to be removed, for a period of 4 hours. The reaction mixture was cooled to room temperature, diluted with 1N NaOH, and filtered through a sintered glass funnel. The filtrate was acidified with 1N HCl, the resulting solids were filtered and dried to get the title product(1.5g) as a beige solid, mp>295° C.

Analysis: Calcd: C 57.64; H 6.22; N 3.20; S 14.65
Found: C 57.48; H 6.19; N 3.25; S 14.73

¹H NMR (DMSO-d₆): δ: 0.82-0.95(6H, m); 1.32-2.06(8H, m); 2.54(1H, d); 3.93(2H, q); 4.70(1H, broad s); 6.23(1H, s); 6.93(1H, d); 7.60(6H, broad s); 7.84(1H, d); 8.30(1H, s)

Synthetic Example 12

Preparation of (7R,9R)-7-butyl-7-ethyl-6,7,8,9-tetrahydro-2-phenyl-1,3-dioxolo[4,5-b]benzothiazepine-5,5-dioxide

The product (0.74g) from Synthetic Example 7 was dissolved in DMF (5 ml). Potassium carbonate (0.50g) and bromochloroethane (0.47g) were added to the reaction mixture and stirred at 110° C for 2 hours. The mixture was filtered through Celite, washed with EtOAc, and the filtrate was dried and concentrated to get an oil. Chromatography on silica gel, using hexane:EtOAc (1:1) as eluant, afforded the title product (0.68g) as a white solid, mp 71-73° C.

Analysis: Calcd. C 65.81; H 6.78; N 3.49; S 7.99
Found: C 65.89; H 6.80; N 3.50; S 8.08

¹H NMR(DMSO-d₆), δ: 0.71-0.85(6H, m); 1.05-2.12(8H, m); 2.49(1H, d); 3.25(2H, q); 3.42(2H, s); 5.91(1H, d); 6.06(1H, s); 7.27-7.41(6H, m)

Synthetic Example 20

Preparation of (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8,9-dimethoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide

(a) 2-Hydroxy-3,4-methoxybenzaldehyde

Aluminum chloride (21.8g) was added, spatula-wise, to an ice-chilled solution of benzoyl chloride (22.1g) and 1,2,3-trimethoxybenzene (25.0g) in 1,2-dichloroethane (250 ml). The reaction mixture was stirred at 0-5° C for 3 hours, then heated to reflux for 2 hours. The reaction mixture was then poured onto ice/concentrated HCl (100 ml) and stirred for 30 minutes, then extracted with diethyl ether. The organic layer was separated, dried and concentrated to get a solid (23.0g). Chromatography on silica gel, using toluene:EtOAc (9:1) as eluant, afforded the title product (18.0g) as a white solid, mp 127-128° C. ¹H NMR consistent with the desired structure.

(b) (±)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8,9-dimethoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide

The product from step (a) was converted to the title product following the procedures used in steps (a) to (f) of Synthetic Example 10. The title product was isolated as a white solid, mp 142-144° C.

Analysis: C 66.16; H 7.48; N 3.35; S 7.68
Found: C 66.03; H 7.53; N 3.28; S 7.77

¹H NMR(DMSO-d₆), δ: 0.64(3H, t); 0.81(3H, t); 0.87-2.08(8H, m); 2.42(1H, d); 3.73(2H, q); 3.75(3H, s); 3.79(3H, s); 5.50(1H, d); 6.05(1H, d); 6.97(1H, d); 7.27-7.41(5H, m)

Synthetic Example 21

Preparation of (3R,5R)-3-butyl-3-ethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepin-4-ol-1,1-dioxide

(a) 2-Hydroxy-4,5-Dimethoxy-4'-fluorobenzophenone

A 1.0 M solution of boron trichloride (142 ml) in dichloromethane was added to 4-fluorobenzoyl chloride (16.8 ml) in benzene (200 ml). Next, 3,4-dimethoxyphenol (20.0g) in benzene (100 ml) was added and the reaction mixture was stirred at room temperature for 2 hours. The mixture was then poured onto ice water and allowed to stir for 15 minutes, then 1N HCl (500 ml) was added and stirred at room temperature for 17 hours. The reaction mixture was extracted with EtOAc, the EtOAc was separated, concentrated and dried to give the title product (41.7g) as an orange solid. ¹H NMR consistent with desired structures.

(b) (3R,5R)-3-butyl-3-ethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepin-4-ol-1,1-dioxide

The product from step (a) was converted to the title product following the procedures used in steps (a) to (o) of Synthetic Example 1 and the procedure used in Synthetic Example 2. The title product was isolated as white solid, mp 170-171° C.

Analysis: C 61.18; H 6.70; N 3.10; S 7.10
Found: C 61.28; H 6.78; N 2.99; S 7.27

41

¹H NMR(DMSO-d₆): δ: 0.75-0.85(6H, m); 1.07-2.04(8H, m); 3.35(2H, q); 3.42(2H, s); 3.81(3H, s); 6.07(1H, s); 6.33(1H, s); 7.22(2H, t); 7.39(1H, s); 7.40-7.50(2H, m); 7.96(1H, s)

Synthetic Examples 22-54

Each of the following examples was prepared by a method analogous to that of Synthetic Example 1, by one of the other exemplified routes or by chemical methods known to those in the art. In all cases, ¹H NMR and elemental analysis were consistent with the proposed structure.

- (22) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-methanol S,S-dioxide, mp 122-123° C
- (23) (3R,5R)-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-7-nitro-5-phenyl-1,4-benzothiazepine 1,1-dioxide 0.40 hydrate, mp 122-123° C
- (24) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-7-(methoxymethyl)-5-phenyl-1,4-benzothiazepine 1,1-dioxide, mp 118-119° C
- (25) (+)-Trans-7-bromo-3-buryl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-ol 1,1-dioxide 0.40 hydrate, mp 137-138° C
- (26) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8,9-trimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide, mp 169-170° C
- (27) (3R,5R)-3-buryl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-7,8-diyldiacetate 1,1-dioxide, mp 79-81° C
- (28) (8R,10R)-8-Buryl-8-ethyl-2,3,7,8,9,10-hexahydro-10-1,4-dioxono(2,3-H)(1,4)-benzothiazepine 6,6-dioxide, mp 82° C
- (29) (3R,5R)-3-buryl-7,8-dimethoxy-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine 1,1-dioxide 0.20 hydrate, mp 110-111° C

42

- (30) (+)-Trans-3-buryl-8-ethoxy-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine 1,1-dioxide, mp 45-54° C
- (31) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-(methylthio)-5-phenyl-1,4-benzothiazepine 1,1-dioxide hydrochloride, mp 194-197° C
- (32) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-isopropoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide hydrochloride, mp 178-181° C
- (33) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-8-carbaldehyde 1,1-dioxide, mp 165-170° C
- (34) 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepine-8-yl aspartate
- (35) 3,3-Diethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide, mp 163-164° C
- (36) 3,3-Diethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-8-methoxy-1,4-benzothiazepine-1,1-dioxide mp 101-103° C
- (37) 3,3-Diethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-1,1-dioxide, mp 132-133° C
- (38) 3,3-Diethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-4,8-diol-1,1-dioxide, mp 225-227° C
- (39) (RS)-3,3-Diethyl-2,3,4,5-tetrahydro-4-hydroxy-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide, mp 205-206° C
- (40) (+)-Trans-3-buryl-8-ethoxy-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-4-ol 1,1-dioxide, mp 149-150° C
- (41) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-isopropoxy-5-phenyl-1,4-benzothiazepine-4-ol 1,1-dioxide, mp 109-115° C
- (42) (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8,9-trimethoxy-5-phenyl-1,4-benzothiazepine-4-ol 1,1-dioxide, mp 84-96° C

(43) (3R,5R)-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-4,7,8-triol-1,1-dioxide, mp 215-220° C

(44) (+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-4,7,8-trimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide, mp 169-187° C

(45) (+)-Trans-3-butyl-3-ethyl-5-phenyl-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepin-4-yl acetate S,S-dioxide, mp 154-156° C

(46) 3,3-Diethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol, 1,1-dioxide, mp 177-178° C

(47) 3,3-Diethyl-2,3,4,5-tetrahydro-7-methoxy-5-phenyl-1,4-benzothiazepin-8-ol, 1,1-dioxide

(48) 3,3-Dibutyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol, 1,1-dioxide

(49) (+)-Trans-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl hydrogen sulfate, mp 196.5-200° C

(50) (+)-Trans-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl dihydrogen phosphate

(51) 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl hydrogen sulfate

(52) 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl dihydrogen phosphate

(53) (+)-Trans-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl aspartate

Biological Assay

In vivo inhibition of bile acid uptake

Inhibiting intestinal bile acid absorption with bile acid sequestrants or surgically with partial ileal bypass surgery is an effective way to decrease plasma LDL cholesterol concentration. Another approach to decreasing bile acid absorption is inhibiting the ileal bile acid active uptake transport system. It has been shown that this inhibition, as measured by the fecal excretion of bile acids results in hypocholesterolemic activity¹.

(1) Lewis, M.C.; Brieddy, L.E.; and Root, C. Effects of 2164U90 on Ileal Bile Acid Absorption and Serum Cholesterol in Rats and Mice. *J. Lipid. Research*, 1995, 36, 1098-1105.

Fecal Excretion of Bile Acids

Male Sprague-Dawley rats weighing 220-260 g were housed in individual cages and fed normal chow. The rats were divided into 6 treatment groups of 10 to 12 rats per group. The rats were dosed by oral gavage (1 mL/100 g body weight) with test compounds as a suspension in 0.5% methylcellulose at 9:00 am and 3:30 pm for two days. The control group received 0.5% methylcellulose. Two hours after the morning dose on day two, the rats were given a trace amount (1.3 nmoles) of 23, 24, 25-⁷⁵Se-homochole acid taurine (⁷⁵SeHCAT) in 1.0 mL saline orally. ⁷⁵SeHCAT, a synthetic gamma emitting bile acid analog which is absorbed by the ileal bile acid uptake system similar to taurocholic acid, has been used clinically as a measure of ileal bile acid absorption^{1,2}. Feces were collected over the 24 hr following ⁷⁵SeHCAT administration. Fecal content of ⁷⁵SeHCAT was determined using a Packard Auto-Gamma 5000 Series gamma-counter. Representative data are tabulated in Table I as the % inhibition of ⁷⁵SeHCAT.

(1) Galatola, G.; Jazrawi, R. P.; Bridges, C.; Joseph, A. E. A. and Northfield, T.

C. Direct Measurement of First-Pass Ileal Clearance of a Bile Acid in Humans. *Gastroenterology*. 1991, 100, 1100-1105.

(2) Ferraris, R.; Galatola, G.; Barlotto, A.; Pellerito, R.; Fracchia, M.; Cortino, F. and De La Pierre, M. Measurement of Bile Acid Half-Life Using [⁷⁵Se]HCAT in Health and Intestinal Diseases. *Dig. Dis. Sci.* 1992, 37, 225-232.

TABLE 1 1% Inhibition of $^{75}\text{SeHCAT}$

45

Compound of Example	Dose (mg/kg)		
	1.0	0.3	0.1
1		51	54
7		65	72
9		80	70
10		53	33
11		72	71
14	56	41	
16	44	34	
18	39	12	
24	49	24	
45	26	6	

In comparison, the most active compound specifically disclosed in International Patent Application No. WO 93/16035 produced a 9% inhibition of $^{75}\text{SeHCAT}$ at 1.0 mg/kg in this assay.

Pharmaceutical Composition Examples

In the following Examples, the active compound can be any compound of formula (I) and/or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof. The active compound is preferably (3R,5R)-3-butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide or one of the compounds of Synthetic Examples 2 to 53.

(I) Tablet compositions

The following compositions A and B can be prepared by wet granulation of ingredients (a) to (e) and (a) to (d) with a solution of povidone, followed by addition of the magnesium stearate and compression.

Composition A

46

	mg/tablet	mg/tablet
(a) Active ingredient	250	250
(b) Lactose B.P.	210	26
(c) Sodium Starch Glycolate	20	12
(d) Povidone B.P.	15	9
(e) Magnesium Stearate	5	3
	500	300

Composition B

	mg/tablet	mg/tablet
(a) Active ingredient	250	250
(b) Lactose	150	-
(c) Avicel PH 101	60	26
(d) Sodium Starch Glycolate	20	12
(e) Povidone B.P.	15	9
(f) Magnesium Stearate	5	3
	500	300

Composition C

	mg/tablet
Active ingredient	100
Lactose	200
Starch	50
Povidone	5
Magnesium Stearate	4
	359

The following compositions D and E can be prepared by direct compression of the admixed ingredients. The lactose used in composition E is of the direct compression type.

Composition D

	mg/tablet
Active ingredient	250
Magnesium Stearate	4
Pregelatinised Starch NF15	146
	400

Composition E

	mg/tablet
Active ingredient	250
Magnesium Stearate	5
Lactose	145
Avicel	100
	500

Composition F (Controlled release composition)

	mg/tablet
(a) Active ingredient	500
(b) Hydroxypropylmethylcellulose (Methocel K4M Premium)	112
(c) Lactose B.P.	53
(d) Povidone B.P.C.	28
(e) Magnesium Stearate	2
	700

The composition can be prepared by wet granulation of ingredients (a) to (c) with a solution of povidone, followed by addition of the magnesium stearate and compression.

Composition G (Enteric-coated tablet)

Enteric-coated tablets of Composition C can be prepared by coating the tablets with 25mg/tablet of an enteric polymer such as cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate, or anionic polymers of methacrylic acid and methacrylic acid methyl ester (Eudragit L). Except for Eudragit L, these polymers should also include 10% (by weight of the quantity of polymer used) of a plasticizer to prevent membrane cracking during application or on storage. Suitable plasticizers include diethyl phthalate, tributyl citrate and triacetin.

Composition H (Enteric-coated controlled release tablet)

Enteric-coated tablets of Composition F can be prepared by coating the tablets with 50mg/tablet of an enteric polymer such as cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate, or anionic polymers of methacrylic

acid and methacrylic acid methyl ester (Eudragit L). Except for Eudragit L, these polymers should also include 10% (by weight of the quantity of polymer used) of a plasticizer to prevent membrane cracking during application or on storage. Suitable plasticizers include diethyl phthalate, tributyl citrate and triacetin.

(ii) Capsule compositionsComposition A

Capsules can be prepared by admixing the ingredients of Composition D above and filling two-part hard gelatin capsules with the resulting mixture. Composition B (*infra*) may be prepared in a similar manner.

Composition B

	mg/capsule
(a) Active ingredient	250
(b) Lactose B.P.	143
(c) Sodium Starch Glycollate	25
(d) Magnesium Stearate	2
	420

Composition C

	mg/capsule
(a) Active ingredient	250
(b) Macrogol 4000 BP	350
	600

Capsules can be prepared by melting the Macrogol 4000 BP, dispersing the active ingredient in the melt and filling two-part hard gelatin capsules therewith.

Composition D

	mg/capsule
Active ingredient	250
Lecithin	100
Arachis Oil	100
	450

49

Capsules can be prepared by dispersing the active ingredient in the lecithin and arachis oil and filling soft, elastic gelatin capsules with the dispersion.

Composition E (Controlled release capsule)

	mg/capsule
(a) Active ingredient	230
(b) Microcrystalline Cellulose	125
(c) Lactose BP	125
(d) Ethyl Cellulose	13
	513

The controlled release capsule composition can be prepared by extruding mixed ingredients (a) to (c) using an extruder, then spheronising and drying the extrudate. The dried pellets are coated with a release controlling membrane (d) and filled into two-part, hard gelatin capsules.

Composition F (Enteric capsule)

	mg/capsule
(a) Active ingredient	250
(b) Microcrystalline Cellulose	125
(c) Lactose BP	125
(d) Cellulose Acetate Phthalate	50
(e) Diethyl Phthalate	5
	555

The enteric capsule composition can be prepared by extruding mixed ingredients (a) to (c) using an extruder, then spheronising and drying the extrudate. The dried pellets are coated with an enteric membrane (d) containing a plasticizer (e) and filled into two-part, hard gelatin capsules.

Composition G (Enteric-coated controlled release capsule)

Enteric capsules of Composition E can be prepared by coating the controlled-release pellets with 50mg/capsule of an enteric polymer such as cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate, or anionic polymers of methacrylic acid and methacrylic acid methyl ester (Eudragit L). Except for Eudragit L, these polymers should also include 10% (by weight of the quantity of polymer used) of a

50

plasticizer to prevent membrane cracking during application or on storage. Suitable plasticizers include diethyl phthalate, tributyl citrate and triacetin.

(iii) Intravenous injection composition

Active ingredient	0.200g
Sterile, pyrogen-free phosphate buffer (pH 9.0) to	10 ml

The active ingredient is dissolved in most of the phosphate buffer at 35-40°C, then made up to volume and filtered through a sterile micropore filter into sterile 10 ml glass vials (Type 1) which are sealed with sterile closures and overseals.

(iv) Intramuscular injection composition

Active ingredient	0.20 g
Benzyl Alcohol	0.10 g
Glycofrol 75	1.45 g
Water for Injection q.s. to	3.00 ml

The active ingredient is dissolved in the glycofrol. The benzyl alcohol is then added and dissolved, and water added to 3 ml. The mixture is then filtered through a sterile micropore filter and sealed in sterile 3 ml glass vials (Type 1).

(v) Syrup composition

Active ingredient	0.25g
Sorbitol Solution	1.50g
Glycerol	1.00g
Sodium Benzoate	0.005g
Flavour	0.0125ml
Purified Water q.s. to	5.0ml

The sodium benzoate is dissolved in a portion of the purified water and the sorbitol solution added. The active ingredient is added and dissolved. The resulting solution is mixed with the glycerol and then made up to the required volume with the purified water.

51

(vi) Suppository composition

	mg/suppository
Active ingredient	250
Hard Fat, BP (Witepsol H15 - Dynamil NoBet)	1770
	2020

One-fifth of the Witepsol H15 is melted in a steam-jacketed pan at 45°C maximum. The active ingredient is sifted through a 200lm sieve and added to the molten base with mixing, using a Silverson fitted with a cutting head, until a smooth dispersion is achieved. Maintaining the mixture at 45°C, the remaining Witepsol H15 is added to the suspension which is stirred to ensure a homogenous mix. The entire suspension is then passed through a 250lm stainless steel screen and, with continuous stirring, allowed to cool to 40°C. At a temperature of 38-40°C, 2.02g aliquots of the mixture are filled into suitable plastic moulds and the suppositories allowed to cool to room temperature.

(vii) Pessary composition

	mg/pessary
Active ingredient (63lm)	250
Anhydrous Dextrose	380
Potato Starch	363
Magnesium Stearate	7
	1000

The above ingredients are mixed directly and pessaries prepared by compression of the resulting mixture.

(viii) Transdermal composition


Active ingredient	200mg
Alcohol USP	0.1ml
Hydroxyethyl cellulose	

The active ingredient and alcohol USP are gelled with hydroxyethyl cellulose and packed in a transdermal device with a surface area of 10 cm².


52

CLAIMS

1. The compounds of the formula (I) :




wherein R¹ is a straight chained C₁₋₆ alkyl group; R² is a straight chained C₁₋₆ alkyl group; R³ is hydrogen or a group OR¹¹ in which R¹¹ is hydrogen, optionally substituted C₁₋₆ alkyl or a C₁₋₆ alkylcarbonyl group; R⁴ is pyridyl or optionally substituted phenyl; R⁵, R⁶, R⁷ and R⁸ are the same or different and each is selected from hydrogen, halogen, cyano, R¹⁵-acylide, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, P(O)(OR¹⁵)₂, OCOR¹⁵, OCF₃, OCN, SCN, NHCN, CH₂OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHSO₂R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pCH₂O)_nR¹⁵, O(CH₂)_pSO₃R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein p is an integer from 1-4, n is an integer from 0-3 and R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from hydrogen and optionally substituted C₁₋₆ alkyl; or R⁶ and R⁷ are linked to form a group



wherein R¹² and R¹³ are as hereinbefore defined and m is 1 or 2; and R⁹ and R¹⁰ are the same or different and each is hydrogen or C₁₋₆ alkyl; with the proviso that when R³ is hydrogen either R⁷ is not hydrogen or at least two of R⁵, R⁶, R⁷ and R⁸ are not hydrogen; and salts, solvates and physiologically functional derivatives thereof.


53

2. The compounds as claimed in claim 1 which are of the formula (II)




wherein R¹ to R¹⁰ are as hereinbefore defined and R^{7a} is selected from halogen, cyano, R¹⁵-acetyl, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, SO₂R¹⁵, P(O)(OR¹⁵)₂, OCO¹⁵, OCF₃, OCN, SCN, HNCN, CH₃OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHSO₂R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pCH₂O)_nR¹⁵, O(CH₂)_pSO₃R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein n, p and R¹² to R¹⁵ are as hereinbefore defined; and salts, solvates or physiologically functional derivatives thereof.

3. The compounds as claimed in claim 1 which are of the formula (III):



wherein R¹-R¹⁰ are as defined in claim 1; and salts, solvates and physiologically functional derivatives thereof.


4. The compounds as claimed in claim 1 which are of the formula (IV)



wherein R¹-R¹⁰ are as defined in claim 1; and salts, solvates and physiologically functional derivatives thereof.

54

5. The compounds as claimed in claim 1 which are of the formula (IVa)



wherein R¹-R¹⁰ are as defined in claim 1; and salts, solvates and physiologically functional derivatives thereof.

6. The compounds as claimed in claim 1 wherein:

R¹ and R² are straight chained C₁₋₆ alkyl;

R³ is hydrogen or hydroxy;

R⁴ is unsubstituted phenyl;

R⁵ is hydrogen;

R⁹ and R¹⁰ are both hydrogen; and either

R⁷ is selected from halogen, hydroxy, C₁₋₆ alkoxy, optionally substituted C₁₋₆ alkyl, -S(O)_nR¹⁵, -OC(O)R¹⁵, and -CH₂OR¹⁵ wherein R¹⁵ is hydrogen or C₁₋₆ alkyl; and

R⁶ and R⁸ are independently selected from hydrogen and those groups listed in the definition of R⁷; or
R⁸ is hydrogen and R⁶ and R⁷ are linked to form a group -O-(CH₂)_m-O- wherein m is 1 or 2;

and salts, solvates, and physiologically functional derivatives thereof

7. A compound according to any of claims 1 to 6 wherein R⁶ and R⁷ are both methoxy;

8. A compound selected from the group consisting of:

(3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

55

(3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;

(3R,5R)-7-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(3R,5R)-7-Bromo-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;

(3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-7,8-diol 1,1-dioxide;

(3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepin-7-ol 1,1-dioxide;

(3R,5R)-3-Butyl-3-ethyl-2,3,4,5-tetrahydro-7-methoxy-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine-4,8-diol;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide;

56

(+)-Trans-2-((3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepin-7-yl)methoxy) ethanol S,S-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-hydroxy-5-phenyl-1,4-benzothiazepine-7-carbaldehyde 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-thiol 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-sulfonic acid 1,1-dioxide;

(7R,9R)-7-Butyl-7-ethyl-6,7,8,9-tetrahydro-9-phenyl-1,3-dioxolo(4,5-H)(1,4)-benzothiazepine 5,5-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8,9-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(3R,5R)-3-butyl-3-ethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepin-4-ol 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine-7-methanol S,S-dioxide;

(3R,5R)-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-7-citro-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(+)-Trans-3-butyl-3-ethyl-2,3,4,5-tetrahydro-8-methoxy-7-(methoxymethyl)-5-phenyl-1,4-benzothiazepine 1,1-dioxide;

(3R,5R)-3-butyl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-7,8-diyl diacetate 1,1-dioxide;

(8R,10R)-8-Butyl-8-ethyl-2,3,7,8,9,10-hexahydro-10-1,4-dioxono(2,3-H)(1,4)-benzothiazepine 6,6-dioxide;


57

- (7R,5R)-3-buryl-7,8-diehoxy-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-buryl-8-ethoxy-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-isopropoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide hydrochloride;
- (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-carbaldehyde 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- 3,3-Diethyl-5-(4-fluorophenyl)-2,3,4,5-tetrahydro-8-methoxy-1,4-benzothiazepine 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-8-methoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-4,8-diol 1,1-dioxide;
- (RS)-3,3-Diethyl-2,3,4,5-tetrahydro-4-hydroxy-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-buryl-8-ethoxy-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;
- (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-8-isopropoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;
- (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-7,8,9-trimethoxy-5-phenyl-1,4-benzothiazepin-4-ol 1,1-dioxide;
- (3R,5R)-3-buryl-3-ethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-4,7,8-triol 1,1-dioxide;


58

- (+)-Trans-3-buryl-3-ethyl-2,3,4,5-tetrahydro-4,7,8-trimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide;
- (+)-Trans-3-buryl-3-ethyl-5-phenyl-2,3,4,5-tetrahydro-7,8-dimethoxy-1,4-benzothiazepin-4-yl acetate S,S-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;
- 3,3-Diethyl-2,3,4,5-tetrahydro-7-methoxy-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;
- 3,3-Diburyl-2,3,4,5-tetrahydro-5-phenyl-1,4-benzothiazepin-8-ol 1,1-dioxide;
- (+)-Trans-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl hydrogen sulfate;
- (+)-Trans-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl dihydrogen phosphate;
- 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl hydrogen sulfate;
- 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl dihydrogen phosphate;
- (+)-Trans-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl aspartate; and
- 3,3-Diethyl-2,3,4,5-tetrahydro-1,1-dioxo-5-phenyl-1,4-benzothiazepin-8-yl aspartate.
9. (3R,5R)-3-Buryl-3-ethyl-2,3,4,5-tetrahydro-7,8-dimethoxy-5-phenyl-1,4-benzothiazepine 1,1-dioxide, or a salt, solvate, or physiologically functional derivative thereof

10. A method of treating a clinical condition in a mammal for which a bile acid uptake inhibitor is indicated which comprises, administering to a mammal an effective bile acid uptake inhibition amount of a compound of the formula (I)




wherein R¹ is a straight chained C₁₋₆ alkyl group; R² is a straight chained C₁₋₆ alkyl group; R³ is hydrogen or a group OR¹¹ in which R¹¹ is hydrogen, optionally substituted C₁₋₆ alkyl or a C₁₋₆ alkylcarbonyl group; R⁴ is pyridyl or optionally substituted phenyl; R⁵, R⁶, R⁷ and R⁸ are the same or different and each is selected from hydrogen, halogen, cyano, R¹⁵-acylide, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, P(O)(OR¹⁵)₂, OCOR¹⁵, OCF₃, OCN, SCN, NHCN, CH₂OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHCO₂R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pCH₂O)_nR¹⁵, O(CH₂)_pSO₂R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein p is an integer from 1-4, n is an integer from 0-3 and R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from hydrogen and optionally substituted C₁₋₆ alkyl; or R⁶ and R⁷ are linked to form a group




wherein R¹² and R¹³ are as hereinbefore defined and m is 1 or 2; and R⁹ and R¹⁰ are the same or different and each is hydrogen or C₁₋₆ alkyl; with the proviso that when R³ is hydrogen either R⁷ is not hydrogen or at least two of R⁵, R⁶, R⁷ and R⁸ are not hydrogen; and salts, solvates and physiologically functional derivatives thereof.

11. A method of treating a hyperlipidemic condition in a mammal which comprises, administering to the mammal an effective hyperlipidemic treatment amount of a compound of formula (I):




wherein R¹ is a straight chained C₁₋₆ alkyl group; R² is a straight chained C₁₋₆ alkyl group; R³ is hydrogen or a group OR¹¹ in which R¹¹ is hydrogen, optionally substituted C₁₋₆ alkyl or a C₁₋₆ alkylcarbonyl group; R⁴ is pyridyl or optionally substituted phenyl; R⁵, R⁶, R⁷ and R⁸ are the same or different and each is selected from hydrogen, halogen, cyano, R¹⁵-acylide, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, P(O)(OR¹⁵)₂, OCOR¹⁵, OCF₃, OCN, SCN, NHCN, CH₂OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHCO₂R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pCH₂O)_nR¹⁵, O(CH₂)_pSO₂R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein p is an integer from 1-4, n is an integer from 0-3 and R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from hydrogen and optionally substituted C₁₋₆ alkyl; or R⁶ and R⁷ are linked to form a group




wherein R¹² and R¹³ are as hereinbefore defined and m is 1 or 2; and R⁹ and R¹⁰ are the same or different and each is hydrogen or C₁₋₆ alkyl; with the proviso that when R³ is hydrogen either R⁷ is not hydrogen or at least two of R⁵, R⁶, R⁷ and R⁸ are not hydrogen; and salts, solvates and physiologically functional derivatives thereof.

12. The method of claim 11 wherein the hyperlipidemic condition is atherosclerosis.
13. A method according to claim 11 or 12 which comprises administering a compound of formula (I) wherein R⁶ and R⁷ are each -OCH₃.

14. A pharmaceutical composition comprising a compound of formula (I):




wherein R¹ is a straight chained C₁₋₆ alkyl group; R² is a straight chained C₁₋₆ alkyl group; R³ is hydrogen or a group OR¹¹ in which R¹¹ is hydrogen, optionally substituted C₁₋₆ alkyl or a C₁₋₆ alkylicarbonyl group; R⁴ is pyridyl or optionally substituted phenyl; R⁵, R⁶, R⁷ and R⁸ are the same or different and each is selected from hydrogen, halogen, cyano, R¹⁵-acetylde, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, PO(OR¹⁵)₂, OCOR¹⁵, OCF₃, OCN, SCN, NHCN, CH₂OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHSO₂R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pNR¹²R¹³, O(CH₂)_pSO₃R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein p is an integer from 1-4, n is an integer from 0-3 and R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from hydrogen and optionally substituted C₁₋₆ alkyl; or R⁶ and R⁷ are linked to form a group




wherein R¹² and R¹³ are as hereinbefore defined and m is 1 or 2; and R⁹ and R¹⁰ are the same or different and each is hydrogen or C₁₋₆ alkyl; with the proviso that when R³ is hydrogen either R⁷ is not hydrogen or at least two of R⁵, R⁶, R⁷ and R⁸ are not hydrogen; or a salt, solvate or physiologically functional derivative thereof, at least one pharmaceutically acceptable carrier, and optionally one or more other physiologically active agents.

15. A pharmaceutical composition according to claim 14 comprising a compound of formula (I) wherein R⁶ and R⁷ are each -OCH₃.

16. A method for the preparation of a compound of formula (I):




wherein R¹ is a straight chained C₁₋₆ alkyl group; R² is a straight chained C₁₋₆ alkyl group; R³ is hydrogen or a group OR¹¹ in which R¹¹ is hydrogen, optionally substituted C₁₋₆ alkyl or a C₁₋₆ alkylicarbonyl group; R⁴ is pyridyl or optionally substituted phenyl; R⁵, R⁶, R⁷ and R⁸ are the same or different and each is selected from hydrogen, halogen, cyano, R¹⁵-acetylde, OR¹⁵, optionally substituted C₁₋₆ alkyl, COR¹⁵, CH(OH)R¹⁵, S(O)_nR¹⁵, PO(OR¹⁵)₂, OCOR¹⁵, OCF₃, OCN, SCN, NHCN, CH₂OR¹⁵, CHO, (CH₂)_pCN, CONR¹²R¹³, (CH₂)_pCO₂R¹⁵, (CH₂)_pNR¹²R¹³, CO₂R¹⁵, NHCOCF₃, NHSO₂R¹⁵, OCH₂OR¹⁵, OCH=CHR¹⁵, O(CH₂)_pNR¹²R¹³, O(CH₂)_pSO₃R¹⁵, O(CH₂)_pNR¹²R¹³ and O(CH₂)_pN⁺R¹²R¹³R¹⁴ wherein p is an integer from 1-4, n is an integer from 0-3 and R¹², R¹³, R¹⁴ and R¹⁵ are independently selected from hydrogen and optionally substituted C₁₋₆ alkyl; or R⁶ and R⁷ are linked to form a group



wherein R¹² and R¹³ are as hereinbefore defined and m is 1 or 2; and R⁹ and R¹⁰ are the same or different and each is hydrogen or C₁₋₆ alkyl; with the proviso that when R³ is hydrogen either R⁷ is not hydrogen or at least two of R⁵, R⁶, R⁷ and R⁸ are not hydrogen; and salts, solvates and physiologically functional derivatives thereof, which comprises

- (a) wherein R³ is hydrogen; by oxidation of a compound of formula (V):



wherein R^1 to R^{10} are as hereinbefore defined and l is 0 or 1; or

- (b) wherein R^3 is OH ; by oxidation of a compound of formula (I) wherein R^2 is hydrogen; and optionally,
 - (c) separating the mixture of isomers so obtained and/or covering the compound of formula (I) so formed to a corresponding salt, solvate, or physiologically functional derivative thereof.
17. A compound of formula (I) according to any one of Claims 1 to 9 or a pharmaceutically acceptable salt, solvate, or physiologically acceptable salt, solvate, or physiologically functional derivative thereof for use in medical therapy.
 18. A compound of formula (I) according to any one of claims 1 to 9 or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof for use in the prophylaxis or treatment of clinical conditions for which a bile acid uptake inhibitor is indicated.
 19. Use of a compound of formula (I) according to any one of Claims 1 to 9 or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof in the manufacture of a medicament for the prophylaxis or treatment of a clinical condition for which a bile uptake inhibitor is indicated.

INTERNATIONAL SEARCH REPORT CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D283/10 C07D417/04 C07D513/04 A61K31/55		Later national Application No. PC/GB 95/01884
According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Extensive data have been considered during the international search (nature of data base used, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO.A.93 16055 (THE WELLCOME FOUNDATION LIMITED) 19 August 1993 cited in the application see the whole document, particularly examples 30-34 and 37-39 --- RESEARCH DISCLOSURE, vol. 354, October 1993 ENSWORTH 68, pages 691-3, ANONYMOUS 'Pharmaceutical compounds' see the whole document ---	1-19
X	WO.A.94 18183 (THE WELLCOME FOUNDATION LIMITED) 18 August 1994 see the whole document, particularly examples 13 and 44 ---	1-19
P, X	--- -/-	1-19
<input checked="" type="checkbox"/>	Further documents are listed in the continuation of box C. Special categories of cited documents:	<input checked="" type="checkbox"/> Patent family members are listed in annex.
* A* document defining the general state of the art which is not considered to be of particular relevance * E* other document published on or after the international filing date but not taken into account in the prior art search * L* document which may show doubts on priority claim(s) or which is used to establish the publication date of another document or other special reason (as specified) * O* document referring to an oral disclosure, use, exhibition or other means * P* document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search		Date of mailing of the international search report
21 November 1995		29.11.95
Name and mailing address of the ISA European Patent Office, P.B. 1011 Patentstr. 2 NL - 2220 HV Rijswijk Tel. (+31-70) 460-2040, Tx. 31 631 epo nl, Fax (+31-70) 460-2016		Authorized officer Allard, H

INTERNATIONAL SEARCH REPORT

Inventor and Application No
PC/GB 95/01884C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT
Category 1 Quotations of documents, with indication, where appropriate, of the relevant passages

Reference to claim No.

P.X WO-A-94 18184 (THE WELLCOME FOUNDATION
LIMITED) 18 August 1994
see the whole document, particularly
example 19

1-19

Form PCT/ISA-118 (Recommendations of search dated July 1993)

page 2 of 2

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inventor and Application No
PC/GB 95/01884

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9316055	19-08-93	AU-B- 3508293	03-09-93
		CA-A- 2117485	19-08-93
		EP-A- 0626952	07-12-94
		FI-A- 943775	16-08-94
		JP-T- 7503724	20-04-95
WO-A-9418183	18-08-94	AU-B- 6007794	29-08-94
		CA-A- 2156184	18-08-94
WO-A-9418184	18-08-94	AU-B- 6008894	29-08-94
		CA-A- 2156183	18-08-94

Form PCT/ISA-118 (patent family member) (July 1993)

INTERNATIONAL SEARCH REPORT		International Application No. PC/GB 95/01884
A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D281/10 C07D417/04 C07D513/04 A61K31/55		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D 000		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where possible, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Class of documents, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO.A.93.16055 (THE WELLCOME FOUNDATION LIMITED) 19 August 1993 cited in the application see the whole document, particularly examples 30-34 and 37-39 ---	1-19
X	RESEARCH DISCLOSURE, vol. 354, October 1993 EHSWORTH GB, pages 691-3, ANONYMOUS 'Pharmaceutical compounds' see the whole document ---	1-19
P.X	WO.A.94.18183 (THE WELLCOME FOUNDATION LIMITED) 18 August 1994 see the whole document, particularly examples 13 and 44 ---	1-19
D. STATEMENT OF RELEVANCE Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patents (family members are listed in annex).		
E. SPECIAL CATEGORIES OF CITED DOCUMENTS: "A" documents defining the general state of the art which is not considered to be of particular relevance "B" documents which are cited in the application but which are not cited in the international search report "C" documents which may throw doubt on priority claim(s) or which are cited to establish the publication date of another document or other special reasons (as specified) "D" documents referred to as oral disclosures, i.e. exhibitions or other means "E" documents which are cited in the international search report but which are not cited in the application		
Date of the actual completion of the international search 21 November 1995		Date of mailing of the international search report 29.11.95
Name and mailing address of the ISA European Patent Office, P.O. Box 1201, 7000 La Haye, The Netherlands NL-2220 HV Delft Tel. (+31-70) 340-3040, Telex 31 01 00 00, Fax (+31-70) 340-3016		Authorized officer Allard, H

INTERNATIONAL SEARCH REPORT		International Application No. PC/GB 95/01884
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Class of documents, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P.X	WO.A.94.18184 (THE WELLCOME FOUNDATION LIMITED) 18 August 1994 see the whole document, particularly example 19 ---	1-19

INTERNATIONAL SEARCH REPORT

Information on parent family members

Inventor and Applicant No.
PC/CG 95/01884

Parent document cited in search report	Publication date	Parent family member(s)	Publication date
WD-A-9316055	19-08-93	AU-B- 3508293 CA-A- 2117485 EP-A- 0626952 FI-A- 943775 JP-T- 7503724	03-09-93 19-08-93 07-12-94 16-08-94 20-04-95
WD-A-9418183	18-08-94	AU-B- 6007794 CA-A- 2156184	29-08-94 18-08-94
WD-A-9418184	18-08-94	AU-B- 6008894 CA-A- 2156183	29-08-94 18-08-94